Smithsonian Institution

Libraries



Alexander Wetmore

1946 *Sixth Secretary* 1953

A. Wetmore

M/HF

Coloured engraved title &
4 plates (1 folding & 1 coloured)





Darton & Cº Scu.

THE

NATURALIST'S

AND

TRAVELLER'S

COMPANION;

BY

JOHN COAKLEY LETTSOM, M.D.

the Third Edition.

London, C. Dilly. 1799.

QH
45
.L5

THE

NATURALIST'S AND TRAVELLER'S

COMPANION.

CONTENTS.

PART the FIRST.

SECT. I.

Page

Method of catching and preserving Insects for Collections — — — 1

SECT. II.

Method of preserving Birds and other Animals. — — — 14

SECT. III.

Directions for bringing over Seeds and Plants from distant countries. — 23

SECT. IV.

Method of analizing Mineral Waters. 41

SECT. V.

Of the Contents of the Air. — — 81

SECT. VI.

Page

Directions for collecting and distinguishing Fossil Substances, including Salts, Earths, Metals and Inflammables. — — 93

PART the SECOND.

SECT. I.

Observations and Queries respecting Learning, Antiquities, religious Rites, polite Arts, &c. — — 177

SECT. II.

Commerce, Manufactures, Arts, Trade, &c. 181

SECT. III.

Meteorological Observations, Food, Way of Living, Animal Œconomy in general, &c. — — — 185

SECT. IV.

Zoology. — — — 188

SECT.

SECT. V.

Botany — — — 194

SECT. VI.

Mineralogy. — — — 198

SECT. VII.

Directions for taking off Impressions of Casts from Medals and Coins. — 200

PRE-

PREFACE.

"HE that enlarges his curiosity after the works of nature," says a celebrated writer, "demonstrably multiplies the inlets to happiness. A man that has formed a habit of turning every new object to his entertainment, finds in these productions an inexhaustible stock of materials upon which he can employ himself, without any temptations to envy or malevolence; faults, perhaps, seldom totally avoided by those, whose judgment is much exercised upon the works of art. He has always a certain prospect of discovering new reasons for adoring the sovereign Author of the universe, and probable hopes of making some discovery of benefit to others, or of profit to himself."

No method appears better calculated to enlarge our knowledge of Natural History, than visiting foreign countries, and carefully attending

attending to the different objects they afford, which more or less delight by their novelty and variety; but our inquiries should not be confined merely to private gratification; there are duties of a more rational nature; to be useful to society by distributing happiness amongst our fellow creatures, is one of the highest and most necessary. The numerous products of nature, their application to the wants, the comforts, and even ornaments of life; the manners, customs, and opinions of mankind; agriculture, manufactures, and commerce; the state of arts, learning, and the laws of different nations, when judiciously investigated, tend to enlarge the human understanding, and to render individuals wiser, better, and happier.

The introduction of the common potatoe, the management of silk-worms, the discovery of jesuits bark, the uses of cochineal, lacca and indigo, are undeniable proofs of the advantages which might be derived from the inquiries of ingenious men. The discovery of another such root as the potatoe, another such article of commerce and apparel as silk, another such remedy as the bark, and such other dying articles as cochineal and indigo, would prove acquisitions of the greatest importance to a trading

ing nation, and render the inquiſitive traveller conſpicuous as a public bleſſing.

Many gentlemen and ſea-faring perſons who go abroad, by their office and ſituation in life, enjoy both time and opportunity for collecting the beſt information on ſuch ſubjects of general utility, eſpecially the natural productions peculiar to the place they viſit or reſide in, which they are induced to overlook, for want of proper directions for diſtinguiſhing and preſerving them, whereby things of value and uſe are loſt to the public, and the time of the traveller leſs beneficially employed.

To promote an application of the time and talents of ſuch perſons to rational and commendable inquiries of this kind, is the deſign of the following directions, which the author thinks himſelf juſtified in recommending, as they principally reſult from experiment and obſervation: Theſe were firſt printed in the year 1772, and the reception from the publick was ſuch as to encourage a ſecond edition in 1774; but this likewiſe being ſoon out of print, a third edition was not long afterwards prepared for the preſs, and ſome of the firſt ſections were printed off ſeveral years ago;

various avocations then intervening, prevented the completion of the remaining ſections; nor would this little performance have been now reſumed, had any publication appeared calculated to preclude the utility of the original plan, which has ſince been conſiderably improved, to make it more deſerving of future encouragement.

In the ſecond part are introduced ſeveral queries and obſervations on natural hiſtory, and upon ſubjects in general, which have not been clearly and ſufficiently aſcertained, and therefore merit the attention of the naturaliſt and traveller.

The principal of thoſe writers who have appeared in different departments of Natural Hiſtory, will be noticed under the particular heads they treat upon.

If perſons who go abroad, or reſide in foreign countries, were acquainted with mathematics and drawing, they would in all probability make their remarks more acceptable, by adding accurate maps of the countries they viſit or reſide in; and by joining to them the drawings of men, their dreſſes, utenſils, weapons, coins, machines, rites, ſacrifices, buildings, temples, idols, and

and antiquities; as well as the curious quadrupeds, birds, reptiles, fiſh, inſects, and ſhells peculiar to each place; with the plants found in thoſe climates, eſpecially ſuch as are employed for food, in commerce, manufactures, phyſic, dying and other purpoſes.

In the drawings and deſcriptions relative to natural hiſtory, it is neceſſary to attend to many circumſtances which are the characteriſtics of each ſpecies of the animal and vegetable creation. In quadrupeds the number and diſpoſition of the teeth; ſhape and poſition of the horns; number of the toes in each foot; ſhape and ſize of the claws and hoofs; ſize of the ears; colour and diſpoſition of the whiſkers; nature and growth of the hair in the fur, mane, and tail; length and uſes of the tail; whether calculated to graſp any object, or to give the animal ſtability; and even the attitude which is characteriſtic of the animal, and ſhews beſt it's marks, ſpots, ſtripes, claws, ears, tail, &c. ought to be expreſſed.

In birds, the ſhape and uſes of the bill, whether notched, ſerrated, or otherwiſe remarkable; number and diſpoſition of the toes, and whether diſtinct, lobated, or palmated;

mated; length of the legs and nakedneſs of the knees; colour of the greater and ſmaller quill feathers, upper and under coverts of the wings; number and colour of the tail feathers, and coverts of the tail; appearance of the vent, belly, breaſt, throat, back, creſt, wattles, carunculæ, ſpurs, &c.; attitude peculiar to the bird, and the difference between males and females, and young and old birds, ſhould be deſcribed.

In tortoiſes and turtles the diſpoſition of the ſhell, and it's compartments; ſhape and number of it's toes; abſence or preſence of the tail, and ſhape of the head, muſt be delineated.

In ſnakes, the ſcales above and below, their number, colour, and figure; form of their heads, and whether they are venomous or not, ſhould be remarked.

In fiſh, it is neceſſary to attend to the proportion of the breadth to the length; form of the head and diſpoſition of the palate and teeth; ſhape and poſition of the mouth; ſize and ſituation of the eyes; coverts of the gill, and rays of it's underpart; ſpines, horns, and protuberances of the head; number, figure, ſize, and colour of

of the fins and tail, with the ſpinoſe and ſoft rays in each; the turn of the lateral line, with the form, colour and diſpoſition of the ſcales.

In inſects, the ſeaſon when each of the different kinds appear ſhould be obſerved; the number, ſubſtance, and particular ſhape of their wings, with the poſition of them when the inſect is at reſt; the ſhape of the antennæ or horns, with the number of joints in each; the form of the head, mouth and eyes, more particularly of the head in beetles, of the mouth in bees, waſps, flies, and gnats, and of the eyes in ſpiders; the number and ſize of the legs; the ſhape of the thighs, feet and claws; the ſtings peculiar to the hymenoptera claſs, and the uſes they are applied to: but the natural hiſtory of inſects ſhould in a peculiar manner engage the traveller's attention, as it is of more conſequence to diſcover the natural hiſtory of one deſtructive or uſeful inſect, than merely to collect and bring over twenty in their perfect ſtate; the former, at the ſame time that it makes the ſcience more entertaining, bids fair to benefit mankind, while the latter ſerves only to fill the cabinets of the curious; he ſhould therefore carefully obſerve the manner in which in-

ſects copulate, and the places where they depoſit their eggs; what food the young larvæ or caterpillars feed upon; if vegetable, whether it be the root, trunk, leaf, flower or fruit; if deſtructive, as they moſtly are, the methods uſed by the natives to deſtroy them; and if uſeful, the means of cultivating them; and what are their natural enemies; the form, attitude, and markings of the caterpillar ſhould be deſcribed; if it has feet, their number, and the particular rings on which they are ſituated; whether it be ſmooth, hairy or ſpinous, and the manner of it's changing into the chryſalis or pupa ſtate, and how long it continues before it arrives to perfection, with the various inſtincts and contrivances they have for avoiding dangers and catching their prey.

In ſhells, not to neglect the number of them belonging to one animal; when ſingle, the turn in the windings, whether to the right or left; the ſtripes, ſpots, bands, knobs, ſpines, furrows and other marks; ſhape of the mouth or opening of the ſhell, lips and beaks: In thoſe that have two or more valves, their equal or different ſizes; the form of the hinges where they are connected, and the number of indentures tallying together; the ſtripes and furrows on the

the outſides, and whether longitudinal or tranſverſal; and the animals inhabiting the ſhells ſhould likewiſe be obſerved and delineated.

In the reſt of the worm tribe, the ſhape, arms, and other parts of the animal ſhould be delineated.

In plants the greateſt accuracy is requiſite, the ſhape of the flower being ſo varied, nice attention is neceſſary to diſtinguiſh it's minute parts; the figure and number of the flower leaves; the form and ſections of the flower-cup; the number and diſpoſition of the duſt veſſels, and of the columnar veſſels ſtanding on the fructification (which are reckoned by botaniſts to be the male and female parts of the flower, and in ſome inſtances are on different plants; in others on the ſame plant, but in different parts of it); the ſhape, ſtructure, and colour of the ſtalk and leaves; the appearance and ſtructure of the roots, and ſuch other circumſtances as characterize the different ſpecies of plants, ought never to be omitted.

The following method of preſerving ſeeds, by John Sneyd, eſq. ſhould have been inſerted, under the directions for bringing over

ſeeds, &c. § III. p. 23, which I have extracted from the Tranſactions of the Society for the Encouragement of Arts, Vol. xvi. p. 265. It is merely packing up ſeeds in abſorbent paper, and ſurrounding the ſame by raiſins, or brown moiſt ſugar; which, by experiment, ſeems to afford that genial moiſture requiſite to preſerve the ſeeds in a ſtate fit for vegetation.

The Naturaliſt ſhould endeavour to keep an accurate journal, wherein all the occurrences, obſervations, places, diſtances, deſcriptions, accounts, informations, and remarks, ſhould regularly and daily be entered, while recent in memory.

It would be adviſable to write on a label fixed to each object, a number correſponding to the notes in the journal; by which means, at any future period, the object itſelf may be clearly aſcertained.

THE

THE

NATURALIST's and TRAVELLER's COMPANION, &c.

PART the FIRST.

SECT. I.

The Method of catching and preſerving Inſects *for Collections.*

——— Ten thouſand different tribes
People the blaze. To ſunny waters ſome
By fatal inſtinct fly.
———Through the green-wood glade
Some love to ſtray; there lodg'd, amus'd and fed,
In the freſh leaf. Luxurious, others make
The meads their choice, and viſit every flower,
And every latent herb. (*a*)

INSECTS in general are known to moſt people, the ſyſtematic diſtinctions but to few; nor have we any Engliſh names for the greateſt part of them. The general denomination of beetles, butterflies, moths, flies, bees, waſps,

(*a*) Thomſon's Seaſons, Summer, l. 246.

and a few other common names, are all that our language ſupplies. It would, therefore, be in vain to enumerate the immenſe variety of genera and ſpecies to any perſon unſkilled in the ſcience of entomology: we may, however, give directions under general names, where to find and how to catch each kind. (*b*)

I. The Coleoptera (*c*), or firſt great claſs of inſects, including beetles, are found in and under the dung (*d*) of animals, eſpecially of cows, horſes, and ſheep: many of them make holes under the dung three or four inches deep; it will therefore be neceſſary to have an iron ſpade to dig them out, when in ſearch of this tribe of inſects.

Some (*e*) are found in rotten and half decayed wood, and under the decayed bark of trees; on the carcaſes (*f*) of animals that have been dead four or five days; on moiſt bones that have been gnawed by dogs or other animals; on flowers having a fœtid ſmell; and on ſeveral kinds of fungous ſubſtances, particularly the

(*b*) Vide Schoeffer. Elementa Entomologica. Curtis's accurate inſtructions for collecting and preſerving inſects, and his introduction to the knowledge of inſects tranſlated from the Fundamenta Entomologiæ of Linnæus. Amæn. Acad. v. 7.

(*c*) Coleoptera, from κολεος, a ſheath, and πτερον, a wing, are ſuch inſects as have cruſtaceous Elytra, or ſhells, which ſhut together, and, form a longitudinal ſuture down to the back of the inſect, as the beetle, Bupreſtris ignita, fig. 1.

(*d*) Scarabæus, *chafer*. Dermeſtes, *leather-eater*. Hiſter, *mimick-beetle*. Staphylinus, *rove-beetle*. (*e*) Lucanus, *ſtag-beetle*. Cerambyx, *capricorn-beetle*. (*f*) Hiſter. Silpha, *carrion-beetle*.

(*g*) Byrrhus,

the rotten and moſt ſtinking: others (*g*) may be found in a morning about the bottoms of perpendicular rocks and ſand banks, and alſo upon the flowers of trees and herbaceous plants.

Many kinds (*h*) may be caught in rivers, lakes, and ſtanding pools, by means of a thread net, with ſmall meſhes, on a round wire hoop, fixed at the end of a long pole.

In the middle of the day, when the ſun ſhines hot, (*i*) ſome are to be ſeen on plants and flowers, blighted trees and ſhrubs; others (*k*) in moiſt meadows are beſt diſcovered at night, by the ſhining light which they emit.

A great variety (*l*) ſit cloſe on the leaves of plants, particularly of the burdock, elecampane, coltsfoot, dock, thiſtle, and the like; or feed on different kinds of tender herbs (*m*).

Numbers (*n*) may be found in houſes, dark cellars, damp pits, caves, and ſubterraneous paſſages, or on umbelliferous flowers (*o*), on the trunks as well as the leaves of trees; in timber-yards, and in the holes of decayed wood.

Some (*p*) inhabit wild commons, the margins of pools, marſhes, and rivulets; and are likewiſe ſeen creeping on flags, reeds, and all kinds of water-plants.

Multitudes (*q*) live under ſtones, moſs, rubbiſh, and wrecks near the ſhores of lakes and

(*g*) Byrrhus, curculio, *weevil*. Bruchus, *ſeed-beetle*. (*h*) Gyrinus, *whirl-beetle*. Dytiſcus, *water-beetle*. (*i*) Coccinella, *lady-fly*. Bupreſtis, *burn-cow*. Chryſomela, *golden honey-beetle*. Cantharis, *ſoft-winged beetle*. Elater, *ſpring-beetle*. Necydalis, *clipt-winged beetle*. (*k*) Lampyris, *glow-worm*. (*l*) Caſſidë, *tortoiſe-beetle*. (*m*) Meloa, *bliſter-beetle*. (*n*) Tenebrio, *ſtinking-beetle*. (*o*) Cerambyx, Ptinus. (*p*) Leptura, *wood-beetle*. Cicindela, *gloſſy beetle*. (*q*) Carabus, *ground-beetle*.

rivers.

rivers. Theſe are found alſo in bogs, marſhes, moiſt places, pits, and holes of the earth, on ſtems of trees; and in an evening they crawl plentifully along path-ways after a ſhower of rain.

Some (*r*) may be diſcovered in the hollow ſtems of decayed umbelliferous plants, and on many ſorts of flowers and fruits.

II. Another claſs (*s*) of inſects are found about (*t*) bake-houſes, corn-mills, in ſhips, and in all places where meal is kept; on graſs (*u*), and all kinds of field herbage. Some (*v*) of theſe frequent rivers, lakes and ſtanding pools.

III. Butterflies and moths make another great diviſion (*w*). In the day, when the ſun is warm, butterflies (*x*) are ſeen on many ſorts of trees, ſhrubs, plants, and flowers. Moths (*y*) may be ſeen in the day-time, ſitting on walls, pales, trunks of trees, in ſhades, out-houſes, dry holes, and crevices *; on fine evenings they fly about the

(*r*) Forficula, *earwig*.

(*s*) Hemiptera, from ημισυ, half, and πτερον, a wing, have their upper wings uſually half cruſtaceous, and half membranaceous, not divided by a longitudinal ſuture, but incumbent on each other, as the Cimex, fig. 2. (*t*) Blatta, *cockroach*. (*u*) Mantis, *camel cricket*. Gryllus, *locuſt*. Fulgora, cicada, *flea-locuſt*. Cimex, *bug*. (*v*) Notonecta, *boat-fly*. Nepa, *water-ſcorpion*.

(*w*) Lepidoptera, from λεπις, a ſcale, and πτερον, a wing, are inſects having four wings, covered with fine ſcales in the form of powder or meal, as in the butterfly, Papilio Antiopa, fig. 3. (*x*) Papilio. *butterfly*. (*y*) Phalæna, *moth*.

* The beſt method to kill moths, when ſtanding, or ſitting on walls

the places they inhabit in the day time. Some (*z*) are ſeen flying in the day-time over the flowers of honey-ſuckles and other plants with tubular flowers. Inſects of this ſpecies ſeldom ſit to feed, but continue vibrating on the wing, while they thruſt the tongue or proboſcis into the flowers.

IV. Inſects of this claſs (*a*) are found in woods (*b*), hedges, meadows, ſand-banks, walls, pales, fruits, and unbelliferous flowers; ſome (*c*) fly about lakes and rivers in the day.

V. The fifth diviſion (*d*) including waſps (*e*), bees, (*f*), &c. may be ſeen about hedges (*g*), ſhrubs, flowers, and fruits. Waſps and bees are the only winged inſects that have any great degree of poiſon in them; they ſhould therefore be taken with a pair of forceps, and handled cautiouſly on account of their ſtings, which are dangerous

walls, pales, &c. is to give them a gentle preſſure on the thorax, or breaſt, with the end of any thing ſmooth, as a tobacco-ſtopper, or tooth-pick caſe: this will deprive them of ſenſe and motion for the preſent moment, and they will fall down on whatever is held under them; a pin ſhould then be ſtruck through the thorax, which ſhould be preſſed between the thumb and finger of the left-hand, till it is felt to give way a little; they may then be ſtuck in the receiving box.

(*z*) Spinx, *hawk-moth*.

(*a*) Neuroptera, from νευρον, a nerve, and πτερον, a wing, have four membranaceous tranſparent naked wings, generally like network, as in the Panorpa Coa, fig. 4. (*b*) Myrmeleon, hemerobius, *pearl-fly*. Raphidia, *camel-fly*. (*c*) Libellula, *dragon-fly*, Ephemera, *may-fly*. Phryganea, *ſpring-fly*.

(*d*) Hymenoptera, from υμην, a membrane, and πτερον, a wing. Inſects with four membranaceous wings, tail furniſhed with a ſting; as in the Tenthredo, fig. 5. (*e*) Veſpa, *waſp*. (*f*) Apis, *bee*. (*g*) Tenthredo, *ſaw-fly*. Sirex, *tailed waſp*. Ichneumon,

dangerous. Some (*h*) of this diviſion have ſtings, but no poiſon, and are to be found on the flowers of umbelliferous plants, when the ſun ſhines hot in the middle of the day; at which time others (*i*) are ſeen on ſand-banks, walls, and pales.

VI. Flies of various kinds conſtitute the next claſs (*k*); they fly about the tops of (*l*) trees, little hills, horſes, cows, ſheep, ditches, dung-hills, and every offenſive object. Some (*m*) are found on all ſorts of flowers, particularly thoſe of a fœtid ſmell. Many (*n*) of theſe are moſt eaſily taken when they begin to feed; for in the middle of the day they are ſo quick and active, that it is almoſt impoſſible to catch them.

VII. The laſt great diviſion (*o*) contains ſcorpions, ſpiders, crabs, lobſters, &c. It is neceſſary only to obſerve here, that all kinds of inſects having no wings, may be preſerved in ſpirits, brandy or rum, except crabs, lobſters, and the like, which may conveniently be preſerved dry.

Under each claſs of inſects, I ſhall relate the

mon, *ichneumon-fly*. Sphex, *ichneumon-waſp*. Veſpa. Apis. (*b*) Mutilla, *naked-bee*. (*i*) Chryſis.

(*k*) Diptera, from δυω two, and πτερον, a wing, are ſuch as have only two wings, and poiſers, as in the fly, fig. 6. (*l*) Oeſtrus, *gad-fly*. Muſca, *fly*. Tabanus, *whame*. Hippoboſca, *horſe-fly*.

(*m*) Tipula. Conops. Aſilus, *waſp-fly*. (*n*) Bombylius, *flower-breeze*.

(*o*) APTERA, from α, without, and πτερον, a wing, inſects having no wings, as the ſpider, fig. 7.

methods


1


2

3

methods of killing them the moſt readily, and with the leaſt pain; as the purſuit of this part of natural hiſtory hath often been branded with cruelty; and however reaſonably the naturaliſt may exculpate himſelf by pleading the propriety of ſubmitting to an evil, which leads to uſeful diſcoveries, yet for wanton cruelty there never can be a juſt pretext.

> —The poor beetle that we tread upon,
> In corp'ral ſufferance finds a pang as great
> As when a giant dies (*p*).

I. The firſt claſs, conſiſting of beetle (coleoptera) are hard-winged. Many kinds fly about in the day, others in the evening, ſome at night only. They may be caught with a gauze net, (pl. 2, fig. 2.) or a pair of forceps covered with gauze (pl. 2, fig. 2). When they are taken, ſtick a pin through the middle of one of the hard wings, and paſs it through the body, as in pl. 1. fig. 1, frontiſpiece. They may be killed inſtantly, by immerſion in hot water, as well as in ſpirit of wine; then ſtick them on a piece of cork, and afterwards carefully place their legs in a creeping poſition, and let them continue expoſed to the air until all the moiſture is evaporated from their bodies. Beetles may alſo be preſerved in ſpirit of wine, brandy or rum, cloſely corked up.

II. Inſects of the ſecond claſs (hemiptera) may be killed in the ſame manner as beetles, and likewiſe by means of a drop of the etherial oil of turpentine applied to the head; or in the manner

(*p*) Shakeſpeare's Meaſure for Meaſure.

manner to be deſcribed under the next claſs for killing moths.

III. The diviſion of butterflies and moths (lepidoptera) as well as all flies with thin membranaceous wings, ſhould be catched with a gauze net, or a pair of gauze forceps: (ſee pl. 1, fig. 2) when taken in the forceps, run a pin through the thorax or ſhoulders, between the fore-wings, as in frontiſpiece, fig. 3. After this is done, take the pin by the head, and remove the forceps, and with the other hand pinch the breaſt of the inſect, in order to deprive it of ſenſation and life: the wings of butterflies ſhould be expanded, and kept ſo, by the preſſure of ſmall ſlips of paper, for a day or two. Moths expand their wings when at reſt, and they will naturally take that poſition.

The larger kinds of theſe inſects will not ſo readily expire by this method, as by ſticking them upon the bottom of a cork exactly fitted to the mouth of a bottle, into which a little ſulphur had been put, and by gradually heating the bottle till an exhalation of the ſulphur takes place, when the inſect inſtantly dies, without injuring its colours or plumage.

Perhaps the moſt eaſy method of killing moths and butterflies is by means of a needle made of ſteel fixed in an ivory handle, which muſt have paſſed hrough the body of the moth; the thumb and forefinger of the left-hand muſt be inſtantly placed on each ſide of the cheſt of the moth, juſt under the wings, when a ſqueeze will entirely deprive the inſect of motion, and probably of ſenſation too for ſome minutes; theſe, however, will return

turn unleſs prevented by the following method.

The point of the needle muſt be paſſed through a ſmall hole in thin braſs, or tin plate, and then held in the flame of a candle for about half a minute, which will effectually deſtroy the life of the inſect.

Although the idea of burning the inſect to death may appear cruel at firſt ſight, yet it ſhould be remembered, that this is not done till after the inſect has been deprived of ſenſation, by preſſure firſt recommended; ſo that, on the whole, it appears to be the moſt eaſy and, ſpeedy method of deſtroying them hitherto practiſed.

It is hardly neceſſary to add, that a braſs or tin plate, for which a piece of card or paſteboard may be ſubſtituted, is uſed to prevent the moth from receiving any injury from the candle.

The beſt method of having the moſt perfect butterflies, is to find out, if poſſible, the larva or caterpillar of each, by examining the plants, ſhrubs or trees they uſually feed upon, or by beating the ſhrubs and trees with long poles, and thereby ſhaking the caterpillars into a ſheet ſpread underneath to receive them; to put them into boxes covered with thin canvas, gauze, or cat-gut (pl. 2, fig. 1) and to feed them with the freſh leaves of the tree or herb on which they are found; when they are full-grown, they will change into the pupa, or chryſalis ſtate, and require then no other care, till they come out perfect butterflies, at which time they may be killed, as before directed. Sometimes theſe in-

fects may be found hanging on walls, pales, and branches of trees, in the chryſalis ſtate.

Moths might likewiſe be procured more perfect, by collecting the caterpillars, and breeding them in the ſame manner as butterflies. As the larvæ or caterpillars cannot be preſerved dry, nor very well kept in ſpirit, it would be ſatisfactory if exact drawings could be made of them while they are alive and perfect. It may be neceſſary to obſerve, that in breeding theſe kinds of inſects, ſome earth ſhould be put into the boxes, as likewiſe ſome rotten wood in the corners, and ſome moſs ſhould be put upon it, which ſhould always be kept damp; becauſe, when the caterpillars change into the pupa, or chryſalis ſtate, ſome go into the earth, and continue under ground for many months before they come out into the moth ſtate; and ſome cover themſelves with a hard ſhell, made up of ſmall pieces of rotten wood. Hence alſo, as many go into the earth, valuable inſects may ſometimes be found by digging after them a foot deep, about the roots of trees, ſhrubs, and plants.

IV. The fourth claſs of inſects (neuroptera) may be killed with ſpirit of wine, oil of turpentine, or by the fumes of ſulphur.

V. Thoſe of the next claſs (hymenoptera) may be killed in the ſame manner. A pin may be run through one of their wing-ſhells and body, as repreſented in pl. 1, fig. 5.

VI. Inſects of the ſixth claſs (diptera) may likewiſe

likewiſe be killed by ſpirit, or by fumes of ſulphur.

VII. Thoſe of the laſt diviſion (aptera) are in general, ſubjects which ſhould be kept in ſpirit.

When in ſearch of inſects, we ſhould have a box ſuitable to carry in the pocket, lined with cork at the bottom and top to ſtick them upon, until they are brought home. If this box be ſtrongly impregnated with camphor, the inſects ſoon become ſtupefied, and are thereby prevented from fluttering and injuring their plumage. Beſides a gauze forceps (pl. II, fig. 7) the collector ſhould have a large muſqueto gauze net, made in the ſhape of a bat fowling-net, (pl. II, fig. 3) and alſo a pin-cuſhion with three or four different ſizes of pins to ſuit the different ſizes of inſects.

In hot climates, inſects of every kind, but particularly the larger, are liable to be eaten by ants and other ſmall inſects, eſpecially before they are perfectly dry; to avoid this, the piece of cork on which our inſects are ſtuck in order to be dried, ſhould be ſuſpended from the cieling of the room, by means of a ſlender ſtring or thread; beſmear this thread with bird-lime, or ſome adheſive ſubſtance, to intercept the rapacious vermin of theſe climes in their paſſage along the thread.

After our inſects are properly dried, they may be placed in the cabinet or boxes where they are to remain: theſe boxes ſhould be kept dry; and alſo made to ſhut very cloſe, to prevent ſmall inſects from deſtroying them; the bottoms of the boxes ſhould be covered with pitch, or green wax, over which paper may be laid; or, which

is better lined with cork, well impregnated with a ſolution of corroſive ſublimate mercury.

The fineſt collections have been ruined by ſmall inſects, and it is impoſſible to have our cabinets too ſecure. Such inſects as are thus attacked may be fumigated with ſulphur, in the manner deſcribed for killing moths; if this prove ineffectual, they may be immerſed in ſpirit of wine, without much injuring their fine plumage or colors; and afterwards let them be ſprinkled about their bodies and inſertions of the wings with the ſolution above-mentioned. But baking the inſects in an oven in the manner to be deſcribed in the next ſection for birds, is the moſt effectual method of extirpating theſe enemies; however the utmoſt caution is requiſite in this proceſs, in regulating the heat of the oven.

Theſe obſervations and directions reſpecting inſects, may, perhaps, be the means of exciting the curioſity of ſome, whoſe inquiries after this part of natural hiſtory will be amply compenſated by the frequent opportunities of enlarging their knowledge, as there is ſcarce any part of the ſurface of this globe, ſcarce a tree, a ſhrub, or a plant, an animal either living or dead, or even the excrements of animals, on which ſome kind of inſect does not depend for its ſubſiſtence and propagation.

——— The flowery leaf
Wants not its ſoft inhabitants. Secure,
Within its winding citadel, the ſtone
Holds multitudes. But chief the foreſt-boughs,
That dance unnumber'd to the playful breeze,
The downy orchard, and the melting pulp
Of mellow fruit, the nameleſs nations feed
Of evaneſcent inſects. Where the pool
Stands mantled o'er with green, inviſible,
Amidſt the floating verdure, millions ſtray.
Each liquid too, whether it pierces, ſoothes,
Inflames, refreſhes, or exalts the taſte,
With various forms abounds. Nor is the ſtream
Of pureſt cryſtal, nor the lucid air,
Though one tranſparent vacancy it ſeems,
Void of their unſeen people———(q).

(q) Thomſon's Seaſons, SUMMER, l. 296.

SECT.

SECT. II.

Method of preſerving BIRDS *and other Animals,*

——— Vitam excoluere per Artes.
VIR. Æn. 6. v. 663.

THE general increaſe of knowledge of late in natural hiſtory, from the attention of individuals to various branches of it, muſt afford no ſmall degree of pleaſure to the ſenſible part of mankind. Whilſt ſuch different reſearches have given entertainment to different diſpoſitions, enlarged the mind, and engaged and diverted it from unprofitable or dangerous purſuits, they have occaſionally given riſe to the moſt uſeful improvements in every department of life, and afforded means of joining utility with elegance.

To promote theſe purpoſes more effectually, a more general knowledge of a good antiſeptic for animal ſubſtances has been much enquired after. Owing to a want of this, many curious animals, and birds particularly, come to our hands in a very imperfect ſtate: ſome from foreign parts entirely miſcarry, and others of the fineſt plumage are devoured by inſects.

To

To promote theſe purpoſes more effectually, a knowledge of the means of preſerving birds and other animals muſt be highly deſirable. The methods made uſe of, by captain Davies, and T. S. Kuckahn, have been publiſhed in the philoſophical Tranſactions (*r*).

The former directs birds in perfect plumage, "to be opened from the upper part of the breaſt "to the vent, with a ſharp knife, or pair of ſciſſars, "the feathers of the breaſt and belly being firſt "carefully laid aſide by the fingers, ſo as not to "hinder the ſkin being eaſily come at. The "ſkin muſt then be carefully looſened from all "the fleſhy parts of the breaſt, body, thighs, "and wings; after this, cut off all the fleſh from "thoſe parts, and take out alſo the entrails and "all the inſide; then, having got a compoſition "of burnt alum, camphor, and cinnamon, of each "an equal quantity, well powdered and mixed "together; ſtrew ſome of this powder lightly "over the whole carcaſe; but ſalt is by no "means to be uſed in this compoſition, as it al-"ways will drop and naſty the plumage in moiſt "weather: pour alſo into the body a ſmall "quantity of camphor diſſolved in rectified ſpi-"rits of wine; after that, fill up the cavity with "fine cotton, or any ſoft woolly ſubſtance, pour-"ing ſome of the aforeſaid ſpirits into the cot-"ton or ſtuffing. Open next the mouth, and "with a pair of ſciſſars take away the tongue, "the roof of the mouth, eyes, brains, and in-"ſide of the head; fill that alſo with the ſame "compoſition; and, having procured eyes as "near the natural ones as poſſible, put them

(*r*) Vol. IX. anno 1770, p. 184, and 302.

into

" into the ſockets by means of a ſmall pair of
" nippers introduced at the mouth. The eyes
" will be beſt made by letting fall ſome drops
" of black ſealing wax on a card of the ſize of
" the natural ones (s); the card muſt be cut
" ſomething larger than the wax, to prevent
" their falling out of the head. Fill the head
" quite full with cotton, pouring ſome of the
" ſpirits down the throat, with ſome of the pow-
" der; a ſmall piece of braſs wire, that has been
" heated in the fire to make it pliable, may be
" put down the throat, being paſſed through one
" of the noſtrils, and faſtened to the breaſt bone,
" to place the head in any attitude you chooſe;
" next fill up the body, where the fleſh has been
" taken away, with cotton and your compoſition;
" and having a fine needle and ſilk, ſew up the
" ſkin, beginning at the breaſt, obſerving, as you
" approach towards the vent, to ſtuff the ſkin as
" tight as it will bear. This will be eaſieſt ac-
" compliſhed by means of a ſmall piece of ſtick
" or ivory, like a ſkewer, till the whole is done:
" then lay your feathers of the breaſt and belly in
" their proper order, and your bird will be com-
" pleted. If you would chuſe to put it into an
" attitude, by introducing a ſmall piece of the
" wire above-mentioned through the ſole of
" each foot up the leg, and into the pinion of

(s) Wax is not a proper ſubſtance for eyes; there are perſons in London, whoſe buſineſs it is to make glaſs eyes of any ſize or colour, at a trifling expence, but as theſe cannot be had in voyages, it would be proper, while the bird is freſh, to take a drawing of the eyes in colours.

" each wing (*u*) it may be diſpoſed of as you " pleaſe."

Inſtead of uſing the ſolution of camphor in ſpirit of wine, Kuckahn reccommends a liquid varniſh, made of two pounds of common or raw turpentine, one pound of camphor, and one pound of ſpirit of turpentine. The camphor is to be broke into very ſmall pieces, and the whole is to be put into a glaſs veſſel, open at top, which is to be placed in a ſand heat, till the ingredients are perfectly diſſolved.

For the dry compound of cinnamon, burnt alum, and camphor, directed in the foregoing method, he ſubſtitutes the two following compoſitions.

Corroſive ſublimate, — —	¼ lb.
Saltpetre prepared or burnt, —	½ lb.
Alum burnt, — —	¼ lb.
Flowers of ſulphur, — —	½ lb.
Muſk (*x*), — — —	¼ lb.
Black pepper, — —	1 lb.
Tobacco ground coarſe. —	1 lb.

(*u*) The late Mr. Leman, who was remarkable for the eaſy attitude of his birds, paſſed a wire ſideways through one wing into the breaſt-bone, the other end of the wire being faſtened into the box incloſing the bird.

(*x*) The muſk renders this compoſition very expenſive for which the ſame quantity of camphor or myrrh might be ſubſtituted.

Mix the whole together, and keep it in a glaſs veſſel ſtopped cloſe. Some of this is to be ſtrewed upon the inſide of the ſkin and cavity of the head, after they have been waſhed with the varniſh.

The other dry compoſition (*y*) is made of equal quantities of tanſy, wormwood, hops, and tobacco, which are to be cut ſmall and mixed together; with this the cavities of the craw and body are to be ſtuffed. He likewiſe directs an artificial breaſt to be made of ſoft wood, and fitted to the proper place, after being moiſtened with the varniſh (*z*).

I thought it might be acceptable to lay before the reader, the above methods, which were practiſed ſome years ſince in this country; but at preſent, a very ſimple one is purſued, which is at the ſame time preferable.

In preſerving all animals, the principal objects ſhould be to remove not only the fleſh of the ſubject but likewiſe as much as poſſible, the bones, which are equally liable to putreſcency, and to invite inſects. In ſhort, after opening the bird,

(*y*) This is entirely uſeleſs, and forms a leſs ſoft and ſmooth ſtuffing than cotton or tow, which on that account are preferable. The reader will obſerve the difficulty and expence of following this complex method recommended by Kuckahn; it is indeed ſurpriſing that his prolix directions ſhould be admitted at large into the Philoſophical Tranſactions.

(*z*) It muſt be almoſt impracticable to proportion an artifical breaſt exactly of the natural ſize and ſhape; cotton or tow anſwers every purpoſe with leſs trouble;

by

by a longitudinal inciſion from the breaſt to the vent, not only the brain and fleſhy parts, but likewiſe all the bones, except thoſe of the legs, and the arch of the ſkull, ſhould be removed: the ſkin is then to be ſtuffed with a ſufficient quantity of cotton or tow, to give the ſubject its natural ſize.

Were the inſide ſurface of the ſkin to be ſprinkled with the dry antiſeptic powder before-mentioned, or with tobacco-duſt, it might probably be ſome preſervative againſt inſects; but it is not at this time an uſual practice, experience having ſhewn that the removal of the fleſh and bones affords the beſt ſecurity.

To exhibit the bird in its proper attitude, there ſhould be ſome ſubſtitute for the back-bone, in order to give firmneſs to the ſubject; for this purpoſe braſs-wire nealed, (that is heated, in the fire, which makes it pliable and retain any form into which it is bent), or a thin piece of lead, may be paſſed down the throat, or through one of the noſtrils, quite to the vent; this ſhould be done, before, or at the time of introducing the cotton or tow. Wires muſt next be introduced through the feet and legs, or between the bone and ſkin of the latter, and faſtened to the longitudinal wire or lead, paſſing from the head to the vent; the other ends of the wire may be fixed in the perch or board, on which the ſubject is to remain.

Having then carefully ſewed up the ſkin, and placed the bird in the attitude we purpoſe it to retain; it ſhould next be expoſed to a dry air, or

to the warmth of an oven heated to a very moderate degree, and it will ſoon acquire a firmneſs which it will afterwards retain; and when this is effected, it ſhould be fixed in a box, ſecured from inſects, glazed in front to exhibit the preparation to view.

When perſons are on long voyages, they have ſeldom leiſure or convenience to go through this proceſs; and indeed were they to effect it, it would be difficult to preſerve the ſubject from inſects.

Under any embarraſſment of this kind, birds and other animals, particularly ſmall ones, may be put into brandy, wine, arrack or firſt runings, without being ſkinned; and the whole proceſs of preparation juſt deſcribed, may be executed when the traveller arrives in port.

Large Sea-fowl have thick ſtrong ſkins, and ſuch may be ſkinned; the tail, claws, head, and feet, are carefully to be preſerved, and the plumage ſtained as little as poſſible with blood.

Baking in a mild heat is not only uſeful in freſh preſervations; but, will alſo be of great

(*a*) Linné deſcribes another method of preſerving fiſh; this is to expoſe them to the air, and when they acquire ſuch a degree of putrefaction, that the ſkin loſes its coheſion to the body of the fiſh, it may be ſlid off almoſt like a glove; the two ſides of this ſkin may then be dried upon paper like a plant, or one of the ſides may be filled with plaſter of Paris, to give the ſubject a due plumpneſs. Vid. Amæn. Acad. Vol. III. A fiſh may be prepared, after it has acquired this degree of putrefaction, by making a longitudinal

ſervice

ſervice to old ones, in deſtroying the eggs of inſects, if they be ſuſpected.

Small quadrupeds, all kinds of reptiles, as ſnakes, lizards, and frogs; fiſh (*a*) of all ſorts and ſmall tortoiſes, with ſea eggs (*b*), and ſea ſtars, may be put into brandy, rum, arrack, or firſt runnings, with the addition of a little alum.

Shells conſtitute an extenſive part of natural hiſtory, and may be collected in great plenty and variety on the ſhores of moſt iſlands and continents. Thoſe which are found with the fiſh in them, are the moſt valuable for the brightneſs of colour, and ſmoothneſs of ſurface, as they loſe that beauty and poliſh, when they have been long expoſed to the ſun. In bivalves, or thoſe having double ſhells, as cockles, oyſters, &c. both the ſhells ſhould always be collected. It is ſufficient in packing up ſhells, to prevent their rubbing againſt each other, which may be effected by means of paper, moſs, ſand, &c. Some of the ſhell-fiſh may be preſerved in ſpirits, as this might prove an uſeful addition to the knowledge of this department of natural hiſtory.

inciſion on the belly, and carefully diſſecting the fleſhy parts from the ſkin, which are but ſlightly attached to it in conſequence of the putreſcency; the ſkin is then to be filled with cotton and tow, as directed for birds, and laſtly to be ſewed up where the inciſion was made.

(*b*) The echini, or ſea eggs, may alſo be dried; they are however liable to be broken.

The

The nefts and eggs of birds, would likewife contribute to increafe the knowledge of natural hiftory, and prove alfo highly ornamental among collections in that branch of zoology.

SECT-

Boxes for conveying Plants by Sea.



The Box with plants shut down with the openings at the ends and front left for fresh air.

The Cask for sowing seeds with the openings defended by Wire.



The Inside of the box shewing the manner of securing the roots of plants surrounded with earth & moss tied with packthread and fastened ways & ways with laths or packthread to keep them steady.



The Box with divisions for sowing different seeds in earth & cut moss.

Boxes for conveying Plants by Sea.



The Inside of the box shewing the manner of securing the roots of plants surrounded with earth & moss tied with packthread and fastened cross & cross with laths or packthread to keep them steady.



The Box with divisions for sowing different seeds in earth & cut moss.

SECT. III.

Directions for bringing over Seeds *and* Plants *from diſtant Countries.*

Nor ev'ry plant on ev'ry ſoil will grow;
The ſallow loves the watry ground, and low;
The marſhes, alders: Nature ſeems t' ordain
The rocky cliff for the wild aſh's reign;
The baleful yew to northern blaſts aſſigns;
To ſhores the myrtles; and to mounts the vines (*c*)

EVERY part of the world has its peculiar productions; and in no objects of natural hiſtory, is the variety more entertaining, and important than horticulture.

The gardens of the curious have already been enriched with many valuable acquiſitions from diſtant countries; but many attempts alſo to introduce ſeveral other plants equally rare, have been unſucceſsfully made, owing to the bad ſtate of the ſeeds or plants when firſt procured, or the method of diſpoſing of them during long

(*c*) Nec vero terræ ferre omnes omnia poſſunt.
Fluminibus ſalices, craſſiſque paludibus alni
Naſcuntur, ſteriles ſaxoſis montibus orni.
Littora myrtetis lætiſſima: denique apertos
Bacchus amat colles; aquilonem et frigora taxi.
VIRG. G. II. 109.

voyages;

voyages; and ſuch accidents as the utmoſt precaution cannot prevent.

For the purpoſe of tranſportation, ripe ſeeds ſhould be choſen, which have been collected in dry weather, and kept dry without expoſing them to ſunſhine; and internally they ſhould be plump, white, and moiſt.

Attention to the ſtate of the ſeeds of Chineſe plants is peculiarly requiſite, as there is reaſon to ſuſpect that more ſeeds from China miſcarry from the art and treachery of the natives, than from the diſtance, or any defect in conveying them; as many ſeeds are brought over, which appear to have been roaſted by the Chineſe, previous to diſpoſing of them, in order to prevent their vegetation, and thereby keep up the demand. To diſcover the healthy ſtate of the ſeeds, ſome of the larger ones may be cut acroſs, and the ſmaller ones bruiſed, and by means of a magnifying glaſs, or even by the naked eye, it may be diſcovered, whether their internal part which contain the ſeminal leaves, appear plump, white and moiſt. If ſo, we may conclude they poſſeſs a vegetating ſtate; but if they are ſhrivelled, inclining to brown or black, and are rancid, they cannot in the leaſt be depended upon.

(a) Seeds thus carefully ſelected, may be preſerved by rolling each in a coat of yellow bees-wax, about half an inch thick; and afterwards a number of theſe, thus prepared, may be put into a chip-box, which is to be filled with melted bees-wax, not made too hot: the outſide of the box may then be waſhed with a ſolution of

of ſublimate mercury (*d*), and kept during the paſſage in a cool airy place. In this manner tea-ſeeds, the ſtones of mangoes, and all hard nuts and leguminous ſeeds in general, may be prepared.

(b) Inſtead of putting ſmall ſeeds in bees-wax, they may be encloſed in paper or cotton which has been firſt ſteeped in melted bees-wax, and then placed in layers in a chip-box, ſome of which may be filled as before with melted bees-wax. Pulpy ſeeds, as thoſe of ſtrawberries, mulberries, arbutuſes, &c. may be ſqueezed together and dried, and then put into the cerate-paper or cotton above-mentioned.

(c) Small ſeeds, may alſo be mixed with a little dry ſand, put into the cerate-paper or cotton, and packed in glaſs-bottles, which are to be well corked, and covered with a bladder or leather (*e*).

(*d*) Sublimate mercury is the moſt effectually diſſolved in the acid of ſea-ſalt; one drop of which will diſſolve one grain of mercury, which will afterwards mix with water. One drachm of ſublimate will be ſufficient for half a pint of water.——— Corroſive ſublimate may likewiſe be diſſolved in a ſaturated ſolution of ſal ammoniac in water, one ounce of which will diſſolve twenty ſcruples of ſublimate.

(*e*) This may be compared with what Dr. Hawkeſworth obſerves in his Voyages. vol. ii. p. 123. "On the 10th I put ſome ſeeds of melons, and other plants, into a ſpot of ground which had been turned up for the purpoſe: they had all been ſealed up, by the perſon of whom they were bought, in ſmall bottles, with roſin, but none of them came up except muſtard; even the cucumbers and melons failed, and Mr. Banks is of opinion that they were ſpoiled by the total excluſion of freſh air." Some ſeeds which I received from North America, encloſed in corked bottles, have ſince been ſown, and germinated. See g.

Theſe bottles may be put into a keg, box, or any other veſſel, filled with four parts of common ſalt, two of ſaltpetre, and one part of ſal ammoniac; or common ſalt alone, if the others cannot be procured, in order to keep the ſeeds cool, and preſerve their vegetative power.

(d) Seeds and nuts, in their pods, may be encloſed in linen or writing-paper, and put into caniſters, earthen-jars, ſnuff-boxes, or glaſs-bottles; the interſtices between the parcels of ſeeds ſhould be filled with whole rice, millet, panic, wheat-bran, or ground Indian-corn well dried. To prevent any injury from inſects, a little camphor, ſulphur, or tobacco, ſhould be put into the top of each canniſter or veſſel, and their covers well ſecured, to exclude the admiſſion of the external air.

(e) Seeds, well dried, may be put into a box, not made too cloſe, upon alternate layers of moſs, in ſuch a manner as to admit the ſeeds to vegetate, or ſhoot their ſmall tendrils into the moſs. In the voyage the box may be hung up at the roof of the cabin, and when the ſhip is at the place of her deſtination, the ſeeds ſhould be put into pots of mould, or boxes, with a little of the moſs alſo about them, on which they had lain.

(f) After every other precaution in tranſporting ſeeds has failed, I have known inſtances of their having been brought from diſtant parts, even from Botany-Bay, and Norfolk-Iſland, by the circuitous voyage of the Eaſt-Indies, in a perfect ſtate of vegetation, which have been merely wrapped in common brown-paper; and as it is the

the eaſieſt and perhaps beſt method of conveying ſeeds, it ſhould never be neglected: American ſeeds are uſually brought over in this manner. The Chineſe paper is generally employed for ſeeds from the Eaſt-Indies, and is probably as good as our common blotting and brown-paper.

Seeds preſerved after the manner (e), (f), as well as that of (d), and likewiſe, for further ſecurity, ſome of the preceding (a), (b), (c), which have been procured in the Eaſt-Indies, may be examined when the ſhip arrives at St. Helena; and ſome of them, which appear in a ſtate of vegetation, ſhould be ſown in the annexed boxes of earth, between the growing plants, as many ſorts as poſſible; ſome of which may ſucceed in caſe of failure of the plants.

More of the ſame ſeeds may be alſo ſown after the ſhip has paſſed the Tropic of Cancer, near the latitude of thirty degrees north. And if very ſmall bits of broken glaſs are mixed with the earth, or thrown plentifully over its ſurface in the boxes, it may prevent mice and rats from burrowing in it, and deſtroying the tender roots of the plants and growing ſeeds.

In whatever method our ſeeds have been preſerved, it ſhould be a conſtant precaution to ſow them as ſoon as they have been expoſed to the external air, otherwiſe they probably will never vegetate.

In order to take up plants or ſhrubs advantageouſly, that are to be tranſported, a mattock and a ſpade ſhould be provided; with the mattock a ſmall trench ſhould be opened round the plant intended to be taken up; the ſpade

ſhould then be put under the root, which muſt be lifted up with a large ball of earth ſurrounding it; and if it ſhould fall off, it muſt be ſupplied with more earth, ſo as to form a ball about the roots of each plant, which muſt be ſurrounded with wet moſs, and carefully tied about with pack-thread to keep the earth about the roots moiſt: loamy earth will continue moiſt the longeſt.

Of each kind the youngeſt plants of ſhrubs and trees that can be found, ſhould be taken; none of them ſhould be above a foot high; as young plants are found by experience to bear removing much better than old ones.

Convenient boxes for the conveyance of plants in long voyages, are made about four feet long, two broad, and two deep; theſe, when half filled with earth, can be conveniently carried by two men holding the rope-handles fixed to their ends. pl. III. fig. 1. 2. 3.

Theſe ſhould be filled about half full of mould, with a few rotten ſticks or leaves at the bottom, and the plants intended to be ſent, planted in it, as ſoon after the ſhip's arrival as poſſible. When the ſhip is about to ſail, and they are ſent on board, hoops are to be nailed to the ſides of the box, in ſuch a manner, that, arching over it, they may cover the higheſt of the plants; ſmall ropes are to be twiſted between theſe, in the form of a net, to prevent the dogs or cats from getting at them, and ſcratching them up, on account of the freſh mould.

For each box ſo hooped and netted, provide a canvas cover, which may, when put on, entirely

protect

BOXES *for conveying* PLANTS *by Sea.*

Fig. 1.



Fig. 2.



Fig. 3.



F. 1. Form of the Box.

2. The same with hoops and loops. a. a. for securing the Canvas.

3. The same netted.

protect it; and, to prevent this cover from being loſt or miſlaid, nail it to one ſide, and fix loops or hooks to the other, by which it may occaſionally be faſtened down.

The late J. Ellis, Eſq. who took ſingular pains to promote the introduction of Exoticks, recommends the conſtruction of a box and caſk agreeable to pl. IV. fig. 1. 2. The firſt ſhews the inſide of the box, and the manner of ſecuring the roots of plants ſurrounded with earth and moſs tied with pack-thread, and faſtened croſs and croſs with laths or packthread to keep them ſteady.

There muſt be a narrow ledge nailed all round the inſide of the box, within ſix inches of the bottom, to faſten laths or packthread, to form a kind of lattice-work, in order to ſecure the plants in their places, as abovementioned.

The caſk fig. 2. is convenient for ſowing of ſeeds, with the openings defended by wire; and as every ſhip has ſpare caſks, one may be readily formed for the conveyance of ſeeds in vegetation or growing plants, agreeable to the annexed engraving. The following is the proportion it ſhould be of; two feet three inches high, two feet bung diameter, and one foot nine inches head diameter; there ſhould be a large opening at the top wired over, the wired part of which might be lifted up at pleaſure, and a lid with hinges to cover it; this may be either circular or ſquare, as will be moſt convenient, the larger the better; and on the upper part of the ſides there may be four or five little openings wired, with doors to each, for the ſake of giving air all round upon ſome occaſions.

Care

Care muſt be taken not to expoſe the young plants to ſtrong ſunſhine: ſometimes, when the lid and doors are open, it may be neceſſary to throw a matt or thin cloth over them, but this muſt depend on the judgment of the perſon who has the care of them; there ſhould be handles fixed to the ſides, to move it with more ſafety.

There ſhould be a layer of wet moſs, of two or three inches deep, at the bottom of the box or caſk; or, if that cannot be got, ſome very rotten wood or decayed leaves, and then freſh loamy earth, about twelve inches deep, both of which will ſink to a foot deep: the wet moſs is intended to retain moiſture, and to keep the earth from drying too ſoon.

The ſurface of the earth ſhould be covered with moſs cut ſmall, which now and then on the voyage ſhould be waſhed in freſh water, and laid on the earth again to keep the ſurface moiſt, and to waſh off mouldineſs or ſaline vapours which may have ſettled on it. When the plants come up, it will be proper to ſave what rain-water can be got, which will encourage their growth, and be of more ſervice than the water drawn out of caſks that have been long on board the ſhip.

Theſe kind of boxes or caſks will be very proper to ſow many ſorts of ſuch ſeeds in, as are ſo difficult to be brought from China, and other parts of the Eaſt-Indies, to Europe in a vegetative ſtate; ſuch as the lechee, mangoes, mangoſteens, pepper, marking-nuts, various ſorts of peaches, roſes, oranges, citrons, lemons, &c.

If the plants ſent from theſe countries were planted in pots or boxes, and kept there a year, they might be brought over with very little hazard; or even if they were firſt tranſplanted from the woods into a garden, till they had formed roots, they might be ſent with much more ſafety.

The captain who takes charge of them, muſt be particularly informed, that the chief danger plants are liable to in ſea-voyages, is occaſioned by the minute particles of ſalt-water with which the air is charged, whenever the waves have white frothy curls upon them; theſe particles fall upon the plants, and quickly evaporating, leave the ſalt behind, which, choaking up the pores, prevents perſpiration, and effectually kills the plant; he therefore ſhould never let the covers be off, except on days when the wind is not ſufficiently high to beat the water up into what the ſeamen call white-caps. He muſt not keep them always ſhut up during the voyage; for if he does, they will mould and periſh by the ſtagnation of the air under the covers: and if at any time, by accident or neceſſity, they ſhould have been expoſed to the wind when the waves have white-caps, he muſt be deſired to water them well with freſh water, ſprinkling all the leaves with it, to waſh off the ſalt-drops which cover them. In this manner plants may be brought from almoſt any diſtance; many come from China every year in a flouriſhing ſtate.

If it be convenient to the captain to give up a ſmall part of the great cabin to the plants, this is certainly by far the beſt ſtation for them; nor are

are they much in the way as the place which ſuits them beſt is cloſe to the ſtern windows; in this caſe they need not be furniſhed with their canvas covers; and they may frequently have air, by opening the windows when the weather is quite moderate.

As the Chineſe ingraft many plants which they introduce into their dwelling houſes, and which may be purchaſed in a healthy ſtate of vegetation, it is a neceſſary caution, not to earth ſuch plants in the paſſage too high with mould, as they are thereby liable to rot, or be otherwiſe injured by ſuch means. The rolling of the ſhip has ſometimes removed the earth, in which they are planted, and many fine collections have been deſtroyed by replacing the earth too high on the ſtems, particularly of ſuch plants as have been engrafted, a practice very general with the Chineſe.

From the frequent inconvenience of the rolling of ſhips, which ſometimes unearths the plants, or otherwiſe injures them, were the boxes or pots to ſtand upon a ſtage, moveable like the mariner's compaſs, it would certainly afford the beſt conveyance: this was prepared at conſiderable expence, in the veſſel commanded by Captain Bligh, to convey the bread-fruit tree, and other vegetables from the South-Seas: the plants he had procured were in high health, when this laudable and princely plan was fruſtrated by a mutiny on board, to be lamented by every man of ſcience and philanthropy. In merchants ſhips, where commerce is the object of the voyage, and where ſpace and convenience

ence can ſeldom if ever be commanded; and therefore the plan of a moveable ſtage cannot here be expected: from a munificent prince the combination of a company, or a ſociety, we muſt look for the ſupport of ſuch an enterprize.

One method of procuring plants from China has been ſucceſsfully adopted by that liberal promoter of horticulture Gilbert Slater Eſq. which may be introduced here. He has ordered every curious drawing of Aſiatic plants by Chineſe artiſts to be procured for him, in the power of ſea-faring gentlemen; theſe drawings are in general not only elegant and accurate; but ſeverally diſtinguiſhed by their names in Chineſe characters: the characters are copied on Chineſe paper, which being thin and almoſt tranſparent, may be accurately traced, and ſuch as refer to rare productions, are diſtributed to captains of ſhips, and other perſons undertaking voyages, with directions to purchaſe ſuch living plants, as correſpond with the character. Sometimes theſe are accompanied with the drawings themſelves. By this circuitous but advantageous mode, in a few years he will acquire a collection of valuable plants, at preſent not known in Europe.

It is much to be regretted, that after the moſt ſcrupulous attention in collecting and tranſporting very valuable plants, they ſhall ultimately be loſt by the tedious proceſs of our Cuſtom-houſe, employed to prevent ſmuggling. I once ſaw ſome fine plants, for which I ſhould have deemed fifty guineas a cheap purchaſe, totally ſpoiled

by being kept there a full month, without any care whatever, not receiving the leaſt ſprinkling of water, whilſt the Thames flowed within twenty yards of their priſon. This does not reſult from any malicious deſign of the officers, who in their vigilance to prevent ſmuggling, thus occaſion the loſs of property, that might not only gratify laudable curioſity, but enrich our poſſeſſions with new products of importance in food, or arts. At this moment I am ſuffering under the injury that a collection of Aſiatic inſects have ſuſtained, of which I do not expect to receive one in a perfect ſtate.

A plan might be adopted, not only to obviate this painful inconvenience, but at the ſame time encreaſe rather than leſſen the revenue, by giving power to the commiſſioners of the cuſtoms to appoint gentlemen of taſte and knowledge in natural hiſtory, whoſe valuation of ſuch ſubjects ſhall govern the duties, which are uſually taken advalorem; and on payment, that the proprietors be permitted to remove their property at pleaſure. That gentlemen might be found to make theſe valuations gratuitouſly I doubt not; and as to the revenue on ſuch ſubjects, I know it would not ſuffer, for many things are, at preſent, not only ſpoiled or totally deſtroyed, but much undervalued by ignorant perſons, many proofs of which I have in my recollection.

When the naturaliſt is in ſearch of vegetable productions, different ſoils and ſituations ſhould be examined; as the ſea, and its ſhores, deep running waters, dikes, marſhes, moors, mountains,

tains, cultivated and barren fields, woods, rocks, &c. afford each their peculiar plants; and wherever any are collected, the particular ſoil and ſituation ſhould be remarked. Sometimes it may prove inconvenient to convey the plants which may be diſcovered, when it would not be ſo to ſend them dried, in the form of a hortus-ſiccus. To do this in the beſt manner, and to make their ſtalks, leaves, &c. lie flat and ſmooth, "The plants ſhould be gathered in a dry day, "after the ſun hath exhaled the dew; taking "particular care to collect them in that ſtate "wherein the generic and ſpecific characters are "moſt conſpicuous; the ſpecimens ſhould be "ſuffered to lie on a table until they become limber, and then they ſhould be laid upon a paſteboard, as much as poſſible in their natural "form, but at the ſame time with a particular "view to their generic and ſpecific characters: "for this purpoſe, it will be adviſeable to ſeparate one of the flowers, and to diſplay the generic character: if the ſpecific character depends upon the flower, or upon the root, a "particular diſplay of that will be likewiſe neceſſary. When the plant is thus diſpoſed upon "the paſteboard, cover it with eight or ten layers "of ſpongy paper, and put it into the preſs (*f*).

(*f*) "The preſs may be prepared by the following directions. "Take two planks of a wood not liable to warp, two inches "thick, eighteen inches long, and twelve inches broad. Get "four male and four female ſcrews, ſuch as are commonly uſed "for ſecuring ſaſh windows. Let the four female ſcrews be let

" Exert only a ſmall degree of preſſure for the " firſt two or three days; then examine it, un- " fold any unnatural plaits, rectify any miſtakes, " and after putting freſh paper over it, ſcrew the " preſs harder. In about three days more, ſe- " parate the plant from the paſteboard, if it is " ſufficiently firm to allow of a change of place; " put it upon a freſh paſteboard, and covering it " with freſh bloſſom-paper, let it remain in the " preſs a few days longer. The preſs ſhould " ſtand in the ſunſhine, or within the influence " of a fire.

" When it is perfectly dry, the uſual method " is to faſten it down with paſte or gum-water (*g*), " on the right-hand inner page of a ſheet of

" into the four corners of one of the planks, and correſpond- " ing holes made through the four corners of the other plank, " for the male ſcrews to paſs through, ſo as to allow the two " planks to be ſcrewed tightly together. It will not be amiſs " to face the bearing of the male ſcrews upon the wood with " iron plates; and if the iron plates went acroſs from corner " to corner of the wood, it would be a good ſecurity againſt " the warping."

This note I have copied from Dr. Withering's Botanical arrangement, and likewiſe the account of drying plants, as his directions are more full and complete than thoſe I formerly introduced into my Naturaliſt's Companion, An. 1772. Where the convenience of this preſs is not at hand, a ſuitable preſſure may be made by weights, or any heavy body.

(*g*) " A ſmall quantity of finely-powdered arſenic or corroſive " ſublimate is uſually mixed with the paſte or gum-water, to " prevent the devaſtations of inſects; but the ſeeds of ſtaves- " acre, finely powdered, will anſwer the ſame purpoſe, without " being liable to corrode, or to change the colour of the more " delicate plants."

" large

" large ſtrong writing-paper. It requires ſome " dexterity to glue the plant neatly down, ſo that " none of the gum or paſte may appear to defile " the paper. When it is quite dry, write upon " the left-hand inner page of the paper, the name " of the plant, the ſpecific character, the place " where, and the time when, it was found; and " any other remarks that may be thought pro- " per. Upon the back of the ſame page, near " the fold of the paper, write the name of the " plant, and it will then be complete for the " cabinet."

" Some people put the dried plants into ſheets " of writing paper, without faſtening them down " at all; and others only faſten them by means of " ſmall ſlips of paper, paſted acroſs the ſtem or " branches.

" Another more expeditious method is, to take " the plants out of the preſs, after the firſt or " ſecond day; let them remain upon the paſte- " board; cover them with five or ſix leaves of " blotting-paper, and iron them with a hot " ſmoothing iron, until they are perfectly dry: " if the iron is too hot it will change the co- " lours; but ſome people, taught by long practice, " will ſucceed very happily. This is quite the " beſt method to treat the orchis, and other ſlimy " mucilaginous plants.

" Another method is, to take the plants when " freſh gathered, and, inſtead of putting them " into the preſs, immediately to faſten them down " to the paper, with ſtrong gum-water; then dip

" a camel-hair pencil into ſpirit-varniſh (*h*), and " varniſh the whole ſurface of the plant two or " three times over. This method ſucceeds very " well with plants that are readily laid flat; " and it preſerves their colours better than any " other."

However beautiful a collection of dried plants may appear in the form of a hortus-ſiccus, yet where duplicates can be procured, it would be acceptable to receive plants both in flower, and in ſeed, dried in a careleſs manner, without nicety in expanding their foliage: by this means ſome of the flowers have been preſerved more entire; and afforded the botaniſt the moſt accurate characters of the plant, which by expoſure to the vapour of hot water, or being ſoaked in lukewarm water itſelf, has expanded, and exhibited the parts of fructification in the moſt perfect ſtate.

The ſeeds of a plant collected when they are ripe, will in a hortus-ſiccus, long retain their vegetative powers, and many of our valuable plants have thus been caſually introduced. It is well known, that the firſt green-tea-tree poſſeſſed by this country, was raiſed by the late John Ellis Eſq. from a ſeed picked out of a canniſter of tea.

(*h*) The ſpirit-varniſh may be made of a quart of highly-rectified ſpirit of wine, five ounces of gum ſandarach, two ounces of maſtich in drops, one ounce of pale gum elemi, and one ounce of oil of ſpike lavender: theſe are to ſtand in a warm place, and be ſhook frequently, to expedite the ſolution of the gum.

The

The impreſſions of plants well taken off upon paper, look very little inferior to the beſt drawings, and may be done with very little trouble. For this purpoſe, ſome printer's ink (*i*), and a pair of printer's balls, ſuch as are uſed for laying the ink on types, are neceſſary. After rubbing theſe balls with a little of the ink, lay the plant betwixt them, and preſs it ſo as to give it ſufficient colour; then take the plant and lay it carefully on a ſheet of paper, and preſs it with the hand, to give the impreſſion of the plant to the paper, which may be afterwards coloured according to nature; a piece of blotting-paper may be placed betwixt the plant and the hand, to prevent the latter from being dirtied by the ink.

An effectual method of ſending a branch of any plant, with the flowers and parts of fructification entire and perfect, is to put them in bottles of brandy, rum, or arrack; but the colour of the plant is often injured.

Corals, corallines, ſponges, &c. inhabitants of the ſea, are found in conſiderable variety near the coaſts of iſlands and continents, particularly in hot climates. Some of theſe are very tender and brittle when dry, and ſhould therefore be carefully packed up in ſand, in order to keep them ſteady, or placed betwixt papers in the manner of an hortus-ſiccus.

In hot climates, the inſects are very rapacious and I have ſeen the fineſt fan-corals, and others of a ſoft texture when firſt taken out of the ſea,

(*i*) Where this cannot be procured, ivory, or lamp-black, ground with boiled linſeed-oil, may be ſubſtituted.

almoſt

almoſt devoured by ants, before they became dry and hard. To prevent injuries of this kind, a little powdered corroſive ſublimate, arſenic, or ſtaveſacre, may be ſprinkled upon theſe productions. Some of the ſmall, and branches of the large ones, might be alſo put into ſpirits, and the parts of them thereby preſerved much more diſtinct; which would ſerve greatly to illuſtrate their natural hiſtory.

SECT. IV.

Method of Analyzing Mineral Waters (a).

Qui autem ad obſervandum adjicit animum, ei etiam, in rebus quæ vulgares videntur, multa obſervatu digna occurrunt.

Bacon, de Augment. Scient.

As many ſprings contain a volatile principle, ſoon liable to be diſſipated, it is neceſſary to make our experiments on the ſpot, in order to diſcover the contents of ſuch waters. Various as theſe contents may appear, the apparatus proper to detect them, may be reduced into a ſmall compaſs.

When we purpoſe to examine any mineral or medicinal ſpring, the ſoil and face of the country ſhould be conſidered, the ſtony or mineral appearance, and particularly whether there are any mineral veins: the degrees of heat of the water ſhould be aſcertained by a thermometer, and its comparative weight to other ſprings in

(*a*) Wallerius in his Hydrologia, Lewis in his notes on Neumann's Chemiſtry, with Rutty, Lucas, Falconer, Monro, Pearſon, Garnett, and other Engliſh writers, have given directions on this ſubject, which ſome late chemical writers on the continent have further elucidated, as Wiegleb, Bergman, Strachling, Struve, Weſtrumb, and Gottling.

the neighbourhood alſo carefully obſerved; after which we may enter on our experiments.

From the ſubſtances contained in mineral waters, they admit of the four following diviſions.

I. ACID AND NEUTRAL SALTS.
II. EARTHLY SALTS.
III. METALLIC SALTS.
IV. SULPHUREOUS.

I. ACID AND NEUTRAL SALTS.

More eminently ſaline, containing the acids, and the compound of acids and alkalis forming neutral ſalts.

1. Fixed air * the only acid found in any quantity pure.
2. Aerated ſoda.
3. Glauber's ſalt.
4. Vitriolated tartar.
5. Saltpetre.
6. Marine ſalt.

†

II. EARTHY SALTS.

1. Epſom ſalt.
2. Magneſian nitre.

* Vitriolic acid is found in extremely ſmall quantity, and that very ſeldom.

† Perhaps Borax, but it is not quite certain that this is a Mineral.

3. Mag-

3. Magneſian muriate.
4. Magneſia, or common Magneſia.
5. Gypſum.
6. Calcarious nitre.
7. Calcarious muriate.
8. Chalk.
9. Alum.

III. METALLIC SALTS.

1. Aerated iron.
2. Vitriolated iron.
3. Vitriolated copper.
4. Vitriolated zinc.
5. Arſenic.

IV. SULPHUREOUS.

1. Hepatic air.
2. Liver of ſulphur.

Water is found mixed with ſome few ſubſtances not here enumerated, as

a. Silicious earth in very fine particles ſometimes gives to water a milkineſs, and

b. Argillaceous earth in ſuch quantity as to give the water the title of ſaponaceous, from its feel and reſemblance.

Some other ſubſtances, as bitumen and naptha, may give Water ſometimes a little character, as they flow out of the ſame ſprings upon the ſurface of ſome waters.

In this view of the ſubſtances contained in mineral waters, we ſee they are almoſt all ſaline,

and as ſome of them are very generally diſtributed through the earth, we find, unfortunately for our inveſtigation of this ſubject, that mineral water generally contains ſeveral of them, ſometimes many, which makes it neceſſary to ſubject portions of the ſame water to a variety of experiments.

Gypſum, calcarious muriate, chalk, Epſom ſalt, muriate of magneſia, and common magneſia, aerated iron, ſea-ſalt, and fixed air are among the moſt common ſubſtances found in mineral waters.

CLASS

CLASS I.

ACID AND NEUTRAL SALTS.

	Synonymous Names.
1. FIXED AIR.	Carbonic acid.
	Carbonaceous acid.
	Cretaceous acid.
	Mephitic acid.
	Aerial acid.
	Atmoſpheric acid.
	Spiritus ſylveſtris.
	Gas ſylveſtre, &c. &c.
	Gas.

This ſubſtance is found in every part of nature that we are acquainted with, in its different ſtates; fixed as it has been called, and aeriform. In our atmoſphere it is found in the proportion of about one in 100. Pits and cavities are frequently filled with it. It exiſts in immenſe quantities combined with calcareous earth, forming chalk and marble; and with many other ſubſtances. It is conſtantly thrown out from animals and vegetables; it is the great product of combuſtion, and formed in ſhort by every proceſs, in which pure air is combined with carbonaceous matter; hence it is no wonder that it is ſo general and ſo frequently found the uniting medium to ſubſtances contained in mineral waters. Mineral waters, which contain this alone, or conſiderably prevailing, have been called Acidulæ; and perhaps hence all mineral waters were formerly termed Acidulæ.

a. Fixed Air is diſcovered in water by its ſparkling, on agitation, and bubbles of air being given out from it.

b. By its pungent and acid taſte.

c. Tincture of litmus added to it, is changed from its blue colour to a red.

d. On its being added to lime water, white clouds are formed. The fixed air combining with the lime forms chalk, which not being ſoluble, is precipitated in a fine white powder.

e. By boiling it, the fixed air is expelled, and the quantity contained in any portion of water may be meaſured, by conveying it through a tube into lime water, and collecting the precipitated chalk; every 30 grains of which contains 18 grains and $\frac{1}{4}$ of fixed air.

2. AERATED SODA.

Mineral Alkali combined with fixed air.

	Synonymous Names.
2. AERATED SODA.	Marine alkali.
	Mineral alkali.
	Foſſil alkali.
	Natron or natrum.
	Soda.
	Baſe of marine ſalt.
	Effervеſcent ſoda.
	Mephite of ſoda.
	Chalk of ſoda.
	Cretaceous ſoda.

This

This ſalt is found in large quantities in Aſia and Africa, and in their mineral waters, and in the waters of ſome parts of Germany; its cryſtals are commonly rhomboidal, extremely ſoluble in water, but when expoſed to the air, fall into powder, it is detected in water by

a. Paper ſtained with the tincture of litmus is changed to a blue colour.

Tincture of litmus is prepared by infuſing litmus, powdered, for a ſhort time, in diſtilled water; with this the paper is ſtained.

b. Paper ſtained with the tincture of brazil wood looſes its red, and is changed to a violet.

The tincture of Brazil wood is made in the ſame manner as the tincture of litmus, but hot water ſhould here be employed.

c. Paper, ſtained yellow by an infuſion of turmeric, is changed to a brown colour.

The infuſion of turmeric ſhould be made with hot water.

d. Upon adding a ſolution of corroſive ſublimate mercury to this water, a white precipitate is produced; if the alkali be in large proportion, this precipitate will be an orange colour. In this inſtance the alkali unites with muriatic acid of the corroſive ſublimate, and the mercury is precipitated.

e. Add a ſmall quantity of Epſom ſalt to this water, it will diſſolve, and no precipitation will be perceived, but give it a very little warmth, and immediately white flakes will be ſeperated: here the mineral alkali combines with the vitriolic acid of the Epſom ſalt, and detaches the magneſia its other conſtituent part, which alſo unites

to

to the fixed air of the mineral alkali. The mineral alkali gives out fixed air enough to redissolve it in the water; but exposing it to a little heat detaches this fixed air, and the magnesia will be precipitated. This magnesia may be further dissolved again with effervescence, by adding any acid.

f. Upon adding a small quantity of sal. ammoniac, the mineral alkali will combine with the marine acid of the sal ammoniac, and detach the volatile alkali, which will fly off, and may be rendered visible by holding a paper moistened with some volatile acid over it, as the nitrous, marine, or acetous. These forming nitrous, marine, or acetous ammoniac, which becomes more visible.

g. By adding a small quantity of a solution of blue vitriol in water, a precipitate of a bright apple green colour will be occasioned by the alkali combining with the vitriolic acid of the blue vitriol, forming Glauber salt, which is soluble in the water, while the fixed air combines with the copper, which being insoluble, is precipitated.

3. GLAUBER SALTS.

Natron combined with Vitriolic Acid.

	Synonymous Names.
3. GLAUBER SALTS.	Sal. mirabile.
	—— catharticus.
	Soda vitriolata.
	Natron vitriolatum.
	Sulphate of soda.

Its crystals are prisms with sixth unequal and striated sides, terminating in dihædrel summits: it has a bitter taste. When exposed to the air, the crystals

cryſtals fall into powder, it is very ſoluble in water. This is found in abundance in ſea-water, and many mineral waters. It is detected by

a. A Solution of muriated ponderous * earth being added to water containing this ſalt, a precipitate, in the form of white clouds will immediately appear, and white precipitate will be gradually depoſited. The ponderous earth leaving the marine acid, will unite with the vitriolic, and form ponderous ſpar, which being inſoluble in water, gives this appearance when precipitated from it, of a white ſoft cloud, peculiar to itſelf: The marine acid combining with the natron, forms marine ſalt, which remains diſſolved in the water.

b. A ſolution of nitre of ſilver † being added, the water firſt becomes of an opal or milky hue, and after ſome time acquires a rediſh caſt, depoſiting at length a ſediment of that colour. Here the vitriolic acid combines with the ſilver, forming ſulphate of ſilver, a ſubſtance almoſt inſoluble in water, therefore precipitated; The nitrous acid, combining with the natron, forms cubic nitre, which remains diſſolved in the water.

c. A ſolution of lead combined with the acetous acid being added, a white percipitate will immediately be formed. The lead combines with the vitriolic acid, forming a ſubſtance inſoluble in water, while the acetous

* Ponderous earth combined with marine acid.

† Silver combined with nitrous acid.

acid, combining with the natron, remains diſſolved in the water.

d. An equal quantity of highly rectified ſpirit of wine being added to water containing this ſalt, the mixture becomes turbid, and by degrees ſmall cryſtals will be formed at the bottom of the veſſel; the ſpirits of wine combining with the water, ſeparates it from the vitriolated tartar, which is precipitated from it in the form of ſmall cryſtals.

4. VITRIOLATED TARTAR.

Kali combined with vitriolic Acid.

	Synonymous Names.
4. VITRIOLATED TARTAR.	Arcanum duplicatum. Sal de duobus. Sal polychreſtum. Kali vitriolatum. Nitrum vitriolatum. Sulphate of potaſh.

It has a bitteriſh taſte, ſuffers no alteration from expoſure to the air; the form of its cryſtals varies much, according to circumſtances under which they are prepared; they are moſtly ſix ſided pyramids, ſometimes ſix ſided priſms, like rock cryſtals: they are ſmall compared with the foregoing.

This ſalt is rarely found in mineral waters.

a. By adding a ſolution of muriated ponderous earth to water containing this ſalt, the ſame appearances take place as deſcribed (49 a.) while the

the muriatic acid combining with the vegetable alkali, forms ſal digeſtivum ſylvii, which remains diſſolved.

b. By the addition of the mercurial ſolution (*c*) yellowiſh clouds will immediately be formed, and in time a precipitate of the ſame colour will be depoſited. The mercury, leaving the nitrous acid, unites with the vitriolic, forming turpith mineral, which is inſoluble in water, though a part of the vitriolic acid and mercury remain diſſolved along with nitre, formed by the nitrous acid uniting with the vegetable alkali.

c. The addition of the ſolution of ſugar of lead in water ſeparates a white precipitate, as mentioned page 49. c. The vitriolic acid unites with the lead, while the acetous acid combines with the vegetable alkali, forming diuretic ſalt, which continues diſſolved.

d. By adding an equal quantity of highly rectified ſpirits of wine, the mixture will become turbid, and by degrees ſmall chryſtals of vitriolated tartar will be formed at the bottom, for reaſons aſſigned page 50. d.

(*c*) Mercury combined with nitrous acid by the aſſiſtance of heat.

5. SALT PETRE.

The compound of nitrous Acid with Vegetable Alkali.

Synonymous Names.

5. SALT PETRE. { Nitre. / Common nitre. / Priſmatic nitre. / Nitre of potaſh.

This is found generally near places inhabited by animals; it is inflammable, its cryſtals are ſix-ſided priſms, terminating in dihedral pyramids, or cut with a ſlope, and frequently channelled.

a. When found in water, it may be detected, by heating the water containing it, then adding a little vitriolic acid to it, this unites with the alkali, forming vitriolated tartar, and detaches the nitrous acid, which being volatile, may be rendered ſenſible, by holding any thing moiſtened with volatile alkali over the ſurface; the vapour of the volatile alkali, uniting with the vapour of the nitrous acid, forms a viſible whitiſh ſmoke, conſiſting of the nitrous ammoniac in fine duſt.

b. By adding an equal quantity of highly rectified ſpirits of wine to this water, it becomes turbid, and this ſalt is ſeparated from it, with the appearances as deſcribed page 50. d.

c. A piece of paper moiſtened with it, and ſuffered to dry, will be found to become touch-paper.

6 COMMON

6. COMMON SALT.

A Compound of Muriatic Acid with Mineral Alkali.

	Synonymous Names.
6. COMMON SALT.	Salt. Sal gemmæ. Sal marinum. Gem ſalt. Sea or marine ſalt. Culinary or kitchen ſalt. Muriate of ſoda.

This ſalt abounds in nature, in vaſt maſſes in the earth; diſſolved in the water of the ſea; and in ſalt lakes; and, therefore, very frequently met with. Its cryſtals are regular tubes; it is detected

a. By adding vitriolic acid to the water containing it, this unites with the mineral alkali of the ſalt, and forms Glauber's ſalt, which remains diſſolved in the water; the muriatic acid is detached, and, being a volatile ſubſtance, may be rendered viſible as the nitrous acid page 52. a. by means of the vapour of volatile alkali held over it; the volatile alkali uniting with the muriatic acid, forms ſal ammoniac, which becomes viſible in this very fine duſt.

b. By adding a ſolution of nitre of ſilver, a white precipitate is immediately formed, which ſome time after will acquire a bluiſh appearance, and this precipitate will not be rediſſolved by the addition

addition of either the nitrous or acetous acids; the ſilver uniting with the muriate acid of the ſalt, forms luna cornea, a ſubſtance ſcarcely at all ſoluble in water; the nitrous acid unites with the mineral alkali of the ſalt, forming cubic nitre, which remains diſſolved in the water.

c. Upon adding the mercurial ſolution, white clouds will immediately appear, and a precipitate of the ſame colour will ſubſide to the bottom.

In this inſtance the muriatic acid of the ſalt unites with the mercury, forming calomel and corroſive ſublimate; the calomel being almoſt inſoluble in water, is precipitated. The corroſive ſublimate remains diſſolved in the water, with the cubic nitre, formed by the nitrous acid and the mineral alkali.

CLASS II.

EARTHY SALTS.

1. Epsom Salt.

Vitriolic Acid combined with Magnesia.

Synonymous Names.

1. EPSOM SALT. { Bitter purging salt. / Magnesian vitriol. / Salt of sedlitz. / Sulphate of magnesia.

It was formerly obtained from wells in the neighbourhood of Epsom, and in the hills that run to the eastward of that place, whose waters contain it abundantly. It has a very bitter taste, is very soluble in water, and is very apt to form short needle-like crystals, when hastily crystalized; otherwise they are quadrangular prisms, terminating in quadrangular pyramids.

A. The magnesian basis may be detected

a. By the addition of lime water, to water containing this salt, which renders it turbid, and a flaky precipitate will be gradually formed. Lime has a stronger affinity to vitriolic acid than magnesia, combines with it and forms gypsum, a small portion of which remains dissolved in the water, the rest is precipitated along with the magnesia.

b. A

b. A ſolution of cauſtic vegetable alkali alſo renders it immediately turbid, and a white flaky precipitate is depoſited. The cauſtic vegetable alkali uniting with the vitriolic acid, forms vitriolated tartar, which remains diſſolved in the water, while magneſia being inſoluble, is precipitated.

c. But if a ſolution of mild vegetable alkali, or vegtable alkali combined with fixed air, be made uſe of, there will be no precipitation; or, if the liquor ſhould grow turbid, it will become clear again on being ſtirred the magneſia being rediſſolved by means of the fixed air that was contained in it. If the liquor be now placed in a warm ſituation, and this fixed air be ſuffered to eſcape, the magneſia will appear in a white precipitate.

d. By addition of the mineral or volatile alkalis, in their cauſtic and mild ſtate, ſimilar effects will be produced, as here mentioned, where vegetable alkali is uſed in its cauſtic and mild taſte.

e. A ſolution of ſoap being added to water containing Epſom ſalt, it becomes turbid, and a number of white flakes are depoſited. The alkali of the ſoap combines with the vitriolic acid, and forms vitriolated tartar, which remains diſſolved in the water, and the unguinous part with the magneſia forming an earthy ſoap.

f. A ſolution of acid of ſugar being added, no precipitation will take place, unleſs ſome calcarious earthy ſalt be mixed with it.

B. The nature of the acid contained may be detected

g. By

g. By adding a ſolution of ponderous earth in marine acid to this water, white clouds will be inſtantly formed, and a white precipitate be gradually depoſited (deſcribed page 49. a.) The magneſia combines with the marine acid, forming muriate of magneſia, which remains diſſolved in the water.

h. Or by adding a ſolution of nitre of ſilver, there will be no precipitate, but the liquor will have an opal appearance, and after ſometime a reddiſh one, depoſiting at laſt a precipitate of this colour. The magneſia, combining with the marine acid, remains diſſolved. The ſilver, combining with the vitriolic acid, forms this precipitate as deſcribed (page 49. b.)

i. The ſolution of nitre of mercury produces yellowiſh clouds, and by degrees a precipitate of the ſame colour, which is turbeth mineral (page 51. b.) The magneſia, uniting with the nitrous acid forms nitrate of magneſia, which remains diſſolved in the water.

k. If a ſolution of ſugar of lead be employed, a white precipitate, vitriol of lead, will be immediately formed by the lead uniting with the vitriolic acid. The magneſia unites with the acetous acid, and remains diſſolved.

100 Grains of vitriol of lead contains 28 grains of vitriolic acid.

l. Highly rectified ſpirits of wine produces a turbid mixture, and by degrees ſmall cryſtals of Epſom ſalt will be formed at the bottom of the veſſel.

m. A ſolution of blue vitriol being employed, produces clouds of a pale green colour, and

after ſome time a precipitate of the ſame colour.

2. MAGNESIAN NITRE.

Nitrous Acid combined with Magneſia.

Synonymous Names.

2. MAGNESIAN NITRE. { Nitre of magneſia. Nitrated magneſia. Nitrate of magneſia.

It is found in the mother water of nitre, and, like it, very rarely found in mineral waters; it has an acrid and very bitter taſte; it imbibes moiſture from the air.

A. a. The magneſian baſis is detected by the ſame means, and with the ſame appearances, as (page 55, a.)

b. By lime water, the nitrous acid uniting with the lime, the magneſia is precipitated in a white flaky appearance. The cauſtic, vegetable, foſſile, or volatile alkali being added, the ſame appearances are produced, the water becomes turbid, and white flakes appear, as deſcribed (page 56, b.) The cauſtic alkali, uniting with the nitrous acid, precipitates the magneſia.

c. So alſo, when employed in their mild ſtate, the ſame appearance will take place, as deſcribed, (page 56, A. c.)

d. With the addition of ſoap, white flakes appear, as deſcribed above, the nitrous acid combining with the alkali of the ſoap, and the magneſia with the oil.

B. f.

B. e. The nature of the acid may be detected, by adding a little vitriolic acid, when the nitrous acid will be detached, as described (page 52, a. and 53, a.)

f. A solution of blue vitriol gives the same appearances as described (page 57, m.)

3. MAGNESIAN MURIATE.

Muriate Acid combined with Magnesia.

Magnesian muriate, or muriate of magnesia, exists in sea water and salt spring waters, but in a more pure state in Epsom and similar waters, is therefore very common. Magnesian muriate has a very bitter hot taste, is very soluble in water, difficult to chrystalize; its chrystals are needle shaped.

A. a. The magnesian basis may be detected in waters containing this salt, by the same means, and with the same appearances, as described (page 58, a. b. c. d.) The lime and alkalis, combining with the muriatic acid, detach the magnesia.

b. By means of a solution of soap in water, the same appearances take place, as described (page 58, d.)

B. c. The acid contained in this water may be detected by the means described (page 59, B. e.) The muriatic acid being volatile, is detached in an invisible vapour which uniting to the vapour of the volatile alkali, forms a whitish smoke, consisting of sal ammoniac.

d. By adding a little of the solution of silver in nitrous acid, white clouds will appear, ac-

quiring in time a bluiſh appearance, as deſcribed (page 53, b.)

e. Or by adding a little of the ſolution of quickſilver in nitrous acid, white clouds will immediately appear, and the ſame appearances take place as deſcribed (page 53, c.)

f. Upon adding a little of the ſolution of blue vitriol, the precipitate of the pale green colour, above deſcribed, will appear.

4. MAGNESIA;

OR

Magneſia combined with fixed air.

Synonymous Names.

4. MAGNESIA.
- Magneſia alba.
- Common magneſia.
- Mild or effervescing magneſia.
- Magneſian earth.
- Aerated magneſia.
- Magneſian chalk.
- Muriatic earth.
- Cretacious magneſia.
- Mephite of magneſia.
- Carbonated, or carbonate of magneſia.
- Magneſian carbonate, &c.

Magneſia is found no where pure, but, combined with other earths, it is found in abundance in nature, in a variety of forms: it is found in the mother water of nitre, and in the other ſtates mentioned in the two preceding pages.

A. a. The magneſia, contained in the water, may be detected by means of a ſolution of cauſtic, vegetable, or foſſile alkali. The cauſtic alkali unites with the fixed air, is rendered mild, and remains diſſolved in the water, while the magneſia is precipitated

b. By ſuſpending a piece of reddened* litmus paper in it, it will looſe the red colour given it by the vinegar, being blue again. And that this is only owing to the magneſia, will be farther proved, by evaporating the water half away, the remaining liquor will be found to poſſeſs no power of changing the colour of the reddened litmus paper, and the fixed air having eſcaped, the magneſia is precipitated in a white powder, and the liquor will be found to be water only.

c. If the piece of paper, that is made uſe of, be ſtained with the tincture of Brazil wood, inſtead of litmus, it will be changed from a red to a violet colour.

d. Or if a ſolution of ſoap be added, as mentioned (page 56, e.) The ſoap is decompoſed, the alkali of the ſoap uniting with the fixed air, and the oil with the magneſia, forming an earthy ſoap.

B. e. The acid baſis will be detected to be fixed air by evaporation, in a cloſe veſſel, as deſcribed (page 46, e.) The fixed air, as it flies off, may be forced into lime water, which it will decompoſe.

* Firſt ſtain the paper with a little infuſion of litmus, then wet it with vinegar.

GYPSUM

5. GYPSUM.

Vitriolic Acid combined with calcarious Earth.

Synonymous Names.

5. GYPSUM.
- Vitriol of lime.
- Calcarious vitriol.
- Selenite.
- Paris plaiſter.
- Sulphate of lime.

This ſalt abounds in nature, and therefore is exceeding commonly found in ſpring water; but ſo ſmall a quantity diſſolves, that it gives very little taſte to water.

A. a. In waters, containing Gypſum, the earthy baſis may be detected by a piece of paper, ſtained with decoction of Brazil wood, being dipped into this water, will tinge it of a violet colour.

b. If a little acid of ſugar be added to water containing it, the water will immediately become turbid, and a white powder be depoſited. The acid of ſugar has a ſtronger attraction to the calcarious earth than the vitriolic acid combined with it, and forms ſaccharine ſelenite, a ſalt extremely difficult of ſolution, therefore precipitated while the vitriolic acid remains diffuſed through the water.

c. A ſolution of the cauſtic vegetable alkali being added, produces white clouds. The calcarious earth is precipitated, in its cauſtic ſtate, by the union of the alkali with its vitriolic acid,

on

on the water being ſtirred. This ſoon diſſolves again; cauſtic calcarious earth or lime being ſoluble in water, but by expoſing it to the air, or adding a little fixed air to it, it becomes chalk, and is again precipitated.

d. Or the mild vegetable alkali being employed, the precipitate will not diſſolve in the water, but fall to the bottom, and will be found to poſſeſs all the properties of common areated calcarious earth, or chalk.

e. So, inſtead of the vegetable alkali, the foſſile or volatile be employed in its cauſtic ſtate, no precipitation is produced, lime being precipitated; but if the mild volatile alkali be made uſe of, a white powder will be precipitated, which will be found to be chalk.

f. A ſolution of ſoap being added to this water, the ſame appearances will take place, as when added to water containing Epſom ſalt. The calcarious earth uniting with the oil, and being precipitated, the vitriolic acid uniting with the vegetable alkali.

B. g. The acid element is diſcovered by the ſolution of muriated ponderous earth; upon adding this, white clouds are immediately produced, as deſcribed (page 49) and a white inſoluble precipitate, which is an artificial ponderous ſpar, will be gradually depoſited at the bottom of the veſſel, as deſcribed (page 49); only here the calcarious earth remains diſſolved in water in union with the marine acid.

h. Upon adding the ſolution of nitre of ſilver, nitre of mercury, ſugar of lead, or blue vitriol, to waters containing Gypſum, the ſame appearances take place as enumerated (page 57).

CALCARIOUS

6. CALCARIOUS NITRE.

Nitrous Acid combined with calcarious Earth.

Synonymous Names.

6. CALCARIOUS NITRE. { Calcarious nitrate. Mother water of nitre. Nitrate of lime.

It is very rarely found in waters; it, has a bitter disagreeable taste, with something of the taste of nitre.

a. By adding the vitriolic acid to water containing this salt, and warming the liquor, a vapour will arise, which may be rendered visible, by placing a stopper, moistened with volatile alkali, over the surface of this liquid. Here the vitriolic acid, combining with the calcarious earth, forms selenite, which, if in any considerable quantity, will be precipitated and form chrystals. The nitrous acid is detached, and, being volatile, will fly off, and at that time becomes visible by the vapour of the volatile alkali, as described (page 52).

b. By the addition of the saccharine acid, the water will immediately become turbid, and a white powder will be deposited at the bottom of the vessel. The saccharine acid, in the same manner, unites with the calcarious earth, and forms calcarious selenite, which, being almost insoluble, is precipitated, while the nitrous is detached, and flies off.

c. A

c. A ſolution of cauſtic, vegetable, foſſil, or volatile alkali, unites with the nitrous acid, forming nitre, which remains diſſolved in the water, and precipitates the calcareous earth in the form of lime, which diſſolves again in the water, and may be detected as related (page 63).

d. And a ſolution of mild alkali, in like manner, unites with the nitrous acid, calcareous earth being precipitated in its mild form, or chalk, as mentioned in the ſame page.

e. On the addition of a ſolution of ſoap, the ſame appearances will take place, as when it is added to water containing Epſom ſalt (page 56. e.)

f. A ſolution of blue vitriol added to water containing this ſalt, clouds of a pale green colour will be formed.

CALCAREOUS MURIATE.

Muriatic Acid combined with calcareous Earth.

	Synonymous Names.
CALCAREOUS MURIATE.	Mother water of marine ſalt.
	Calcareous marine ſalt.
	Fixed ſal ammoniac.
	Muriate of lime.
	Murias calcareous.

It is found wherever ſea ſalt is met with, particularly in ſea water; chalk often contains it; it has a ſalt and very diſagreeable bitter taſte. It cryſ-

talizes in prifms, with four ftriated faces, terminated with fharp-pointed pyramids. It deliquefces immediately on expofure to the air.

A. To water containing this falt, to detect the earthy bafis,

a. Add a little folution of the acid of fugar, the mixture becomes immediately turbid; the acid of fugar, uniting with the calcareous earth, forms a compound fcarcely foluble in water, which is precipitated, leaving the muriatic acid in the water, as defcribed (page 62, b.)

b. Add a little of the folution of either of the alkalis, in a cauftic ftate, white clouds will immediately form, the alkali uniting with the muriatic acid, the calcareous earth is precipitated; but here, on the liquor being ftirred, the precipitate difappears, for the calcareous earth being in its cauftic ftate (Lime) on being mixed with the water, diffolves in it; but if this liquor be expofed to the air, or if fixed air be thrown into it, this lime becomes chalk, which will fall to the bottom.

c. If the folutions of the alkalis be employed in their mild ftate, or combined with fixed air, white clouds will immediately form, as above, which will difappear again, on ftirring the liquor, but on another principle, there being a double affinity; for while the alkali combines with the muriatic acid, the fixed air, which was combined with the alkali, unites with the calcareous earth, and forms chalk; but here the fixed air is detached in fuch abundance, as alfo to diffolve the chalk fo formed in the water; now warm the water, the fuperabundant fixed

fixed air will fly off, and the chalk be precipitated.

d. By adding a ſolution of ſoap, the ſame appearances will take place, as in water containing Epſom ſalt, deſcribed (page 56, e.)

By adding a ſolution of ſilver in nitrous acid, or of mercury in nitrous acid, or of lead in acetous acid, or of blue vitriol, or by adding ſpirit of wine, the ſame appearances will be produced as upon adding theſe ſubſtances to water containing Epſom ſalts, deſcribed (page 57, h. i. k. l. m.)

B. e. The acid may be detected by the ſame means, and with the ſame appearances as deſcribed (page 53 and 54, a. b. c.)

CHALK.

Calcareous Earth combined with fixed Air.

	Synonymous Names.
CHALK.	Creta.
	Cretaceous earth, &c. or Calcareous earth.
	Calcareous ſpar.
	Calcareous carbonate.
	Carbonate of lime.
	Aërated lime.

This is found in immenſe quantities in moſt parts of the earth; at leaſt the ſuperficial parts; of courſe is found diſſolved in almoſt all ſpring waters.

A. a. Upon placing a bit of reddened litmus paper in water containing it, the red colour will gradually diſappear and the blue return; but if a portion of the water be firſt evaporated, it will become turbid, the ſuperabundant fixed air, which keeps the chalk in ſolution, will fly off, and the chalk be depoſited: now place the litmus paper in it, and no alteration of colour will take place. The calcareous earth not being in ſufficient quantity in the water to combine with the vinegar, that gives this red colour to the litmus paper.

b. Upon putting into it a piece of Brazil wood paper, the red colour will be by degrees changed to a violet; but if a portion of the water be evaporated, and the remainder be ſuffered to ſettle as above, no alteration will happen, unleſs the water contain alſo Gypſum, or ſome alkali.

c. On adding the acid of ſugar to this water a precipitate takes place, as deſcribed (page 62. b.) it combines with the calcareous earth, and falls to the bottom, while the fixed air remains diſſolved in the water.

d. A ſolution of ſoap being added to this water, the ſame appearances take place as deſcribed (page 56. e.) The lime and oil forming an earthy ſoap, while the fixed air unites to the alkali.

B. e. The fixed air contained, if in any quantity, may be diſcovered by the method deſcribed (page 46. d. e.) or by adding a little vitriolic acid to it before the diſtillation. The vitriolic acid unites with the lime, and forms Gypſum, which remains diſſolved in the water, while the fixed air is detached.

ALUM.

ALUM.

Vitriolic Acid combined with argillaceous Earth.

Synonymous Names.

ALUM. { Argillaceous vitriol. / Aluminous ſulphate. / Sulphate of alumine.

Alum has a very aſtringent and acid taſte; its cryſtals are of the figure of an octohedron; it is very rarely found in mineral waters.

A. a. The earthy baſis may be detected by the addition of lime water, which occaſions a turbid appearance, and a flaky precipitate will be gradually formed, which is the earth of alum. The vitriolic acid, uniting with the calcareous earth of the lime water, forms Gypſum, which is more ſoluble in water.

b. A ſolution of alkali, either cauſtic or mild, vegetable, foſſil, or volatile, added to aluminous water, produces a white flaky appearance. The alkali combines in the ſame manner with the vitriolic acid, and precipitates the earth of alum, either alone or combined with fixed air, leaving vitriolated tartar, Glauber's ſalts, or volatile ammoniac, according to the alkali made uſe of, diſſolved in the water.

c. By adding a ſolution of ſoap, the aluminous water becomes turbid, and depoſites a number of white flakes, the alkali of the ſoap uniting

ing

ing with the vitriolic acid, and the unguinous part with the earth, forming an earthy ſoap.

B. d. The vitriolic acid may be detected by the addition of a ſolution of muriated ponderous earth, white clouds will be inſtantly formed, and a white inſoluble precipitate, which is an artificial ponderous ſpar, will be gradually depoſited at the bottom of the veſſel, as related in other inſtances, where vitriolic acid conſtitutes one of the elements, (page 49, a.)

e. By adding the ſolution of nitre of ſilver, nitre of mercury, or ſugar of lead, to water containing alum, the ſame appearances will take place as enumerated (page 49, b. page 57, h. i. k.) the aluminous earth uniting with the acids, in the ſame manner as the magneſia in thoſe inſtances.

f. A little ſolution of blue vitriol being added, a green precipitate will be formed, with a ſmall tinge of blue in it.

CLASS III.

Metallic Substances.

AËRATED IRON,

Iron combined with fixed Air.

AËRATED IRON. Synonymous Names.
- Aperient ſaffron of mars.
- Ruſt of iron.
- Martial chalk.
- Martial mephite.
- Carbonate of iron.

Both iron and fixed air being found in almoſt every ſituation, and readily uniting together, and, eſpecially where the fixed air is in any quantity, readily diſſolving in water; it is no wonder that this is among the moſt common ſubſtances met with in ſprings. Waters containing it are called chalybeate, or ferruginous waters; and, from their ſtrengthening medicinal qualities, are very generally uſed. They have ſomewhat of a ſtyptic or irony taſte, mixed with ſome pungency, from the ſuperabundant fixed air they contain.

This water is very eaſily known.

A. a. In being expoſed to the air, or heated, it precipitates a brown powder, which is calx of iron. The fixed air being decompoſed; the water will have a rainbow-coloured pellicle on it.

(1) b. On adding the tincture of galls, a purple-coloured precipitate will appear, which, if in any quantity, will be almoſt black; but if the fixed air be firſt ſuffered to evaporate, and the iron to precipitate as above, the liquor will effect no change with the tincture of galls.

(2) c. On adding the ſmalleſt quantity of Pruſſian lixivium, a beautiful blue colour will be precipitated. The Pruſſic acid, uniting with the iron, forms Pruſſian blue; neither will this happen, if the precipitation a. be ſuffered to take place firſt.

d. On adding a little ſolution of ſoap, the water will become turbid, and white flakes will be precipitated to the bottom of the veſſel; the alkali of the ſoap uniting to the fixed air, and the iron to the oil, forming a metallic ſoap.

e. On adding a ſolution of volatile liver of ſulphur, a precipitate of a very dark green colour is formed. The ſulphur combining with the iron, becomes inſoluble in the water.

B. f. The fixed air may be detected as deſcribed (page 46, a. e.)

(1) *Tincture of galls is made by infuſing powdered galls in ſpirits of wine.*

(2) *Pruſſian lixivium is formed by digeſting Pruſſian blue with caustic vegetable alkali and water, and then adding a little diſtilled vinegar to neutralize any ſuperabundant alkali, if alkali ſhould prevail.*

VITRI-

VITRIOLATED IRON.

Iron combined with Vitriolic Acid

VITRIOLATED IRON.

Synonymous Names.

- Sal Martis.
- Copperas, or green copperas.
- Vitriolum viride, green vitriol.
- Martial vitriol.
- Vitriol of iron.
- Sulphate of iron.

Although this ſalt is formed from the decompoſition of martial pyrites, which decompoſe very readily, and are extremely common; and is formed alſo by other natural proceſſes; and although this ſalt is by no means uncommonly found ſolid, in it's native ſtate, yet it is very rarely met with in mineral waters. Its chryſtals are of a green colour.

It is detected nearly as the former.

A. a. By expoſing it to the air it will not ſo ſoon, or ſo perfectly, depoſite the iron in a brown precipitate as the former page 71. The vitriolic acid is not ſo readily decompoſed; it will have the ſame rainbow coloured pelicle on it.

b. On adding the tincture of galls, the ſame purple colour will appear as deſcribed p. 72,

c. On adding the Pruſſian lixivium the ſame colour will be formed, as page 72, and that although it has been firſt heated.

d. On adding a ſolution of cauſtic alkali the liquor will become yellow, and, after ſome time,

depoſite an ochry ſediment. The alkali, uniting with the vitriolic acid, depoſites the iron, which unites with a portion of pure air, forming a calx of iron.

e. On adding a ſolution of ſoap, the ſame appearances take place as deſcribed (page 72, d.)

f. On adding a ſolution of liver of ſulphur, a black precipitate will immediately appear: if to this black precipitate a few drops of vitriolic acid be added, it will be perfectly rediſſolved as before.

g. If water be added, impregnated with hepatic air, the ſame effect will be produced.

B. h. The nature of the acid may be more perfectly determined, by adding a ſolution of muriated ponderous earth, when white clouds will inſtantly be formed. The ponderous earth, uniting with the vitriolic acid, forms baroſelenite, which is precipitated, while the iron, combining with the muriatic acid, remains diſſolved in the water; for, upon adding the Pruſſian lixivium, or tincture of galls, the blue or purple precipitates will appear,

i. By adding a ſolution of nitre of mercury, a beautiful yellow precipitate will be obtained. The vitriolic acid uniting with the mercury forms turpeth mineral, while the nitrous acid, uniting with the iron, remains diſſolved in the water, as may be known by adding the tincture of galls, or Pruſſian lixivium.

VITRIOLATED COPPER.

Copper combined with Vitriolic Acid.

	Synonymous Names.
VITRIOLATED COPPER.	Blue vitriol.
	Blue copperas.
	Vitriol of cyprus.
	Vitriol of copper.
	Vitriol of venus.
	Sulphate of copper.

Its chryſtals are of a beautiful blue colour, of an oblong rhomboidal ſhape, the taſte is ſtyptic, even cauſtic. This is often found in mineral waters, in the neighbourhood of copper mines.

A. a. It is detected. By adding lime water, when a pale green precipitate will appear, which, upon pouring in more lime water, is rediſſolved, the liquor becoming a ſapphire blue.

b. By adding the cauſtic volatile alkali, the water alſo becomes of a ſapphire blue colour. The volatile alkali uniting with the copper, forms cuprum ammoniacum, which will remain diſſolved in the water, unleſs it be evaporated.

c. But if the mild volatile, mineral, or vegetable alkali be employed, green clouds will be formed, and in time a precipitate of that colour will fall to the bottom. Theſe alkalis unite with the vitriolic acid, while the copper unites with the fixed air of the alkalis, forming what is called mountain copper, which is precipitated; but if much mild volatile alkali be added, it

will rediſſolve the precipitate, and form cuprum ammoniacum.

d. Upon adding a ſolution of Epſom ſalt, or of nitrous or muriated magneſia, the precipitate will be of a pale green colour.

e. Upon adding a ſolution of nitrated or muriated calcarious earth, the precipitate will be ſtill of a more pale green colour.

f. On adding a ſolution of Alum, a green precipitate will be formed, with a tinge of blue in it. In theſe three inſtances, the earths unite with the acid, and the copper is precipitated.

g. If a ſolution of arſenic in water be added, clouds of yellowiſh green colour will be formed, and a ſubſtance of the ſame colour will be precipitated. This is called ſcheel's green.

h. On adding a ſolution of volatile liver of ſulphur, clouds of a blackiſh brown hue will be formed, and a precipitate of the ſame colour fall to the bottom;

i. Or if water, impregnated with hepatic air, be added, the ſame effect will take place.

k. On adding a little of the Pruſſian lixivium, a precipitate of a brown colour will be depoſited.

B. 1. The nature of the acid may be detected by adding a ſolution of muriated ponderous earth, nitre of ſilver, and the other means mentioned (page 49, a. b. &c.)

VITRIOLATED ZINC.

Zinc combined with vitriolic Acid.

VITRIOLATED ZINC.

Synonymous Names.
- White vitriol.
- White copperas.
- Sal vitrioli.
- Vitriol of zinc.
- Goſlard vitriol.
- Sulphate of Zinc.

This is ſometimes found in waters near the mines of zinc; it has a ſtyptic taſte; its chryſtals are colourleſs, and generally tetrahedral priſms, terminating in pyramids.

a. By adding lime water pearl coloured clouds will appear, and a precipitate of this kind will fall to the bottom.

b. Vegetable alkali, mild or cauſtic, being added, a precipitate will appear of a dirty yellowiſh pearl colour.

c. Upon adding cauſtic volatile alkali, a pale orange coloured preciptate will be formed.

d. But if the volatile alkali be employed in its mild ſtate, a precipitate, nearly of a white colour will be formed. Theſe ſeveral ſubſtances unite with the vitriolic acid, and remain diſſolved in the water, whilſt the zinc is precipitated in the form of a calx.

e. Upon adding the Pruſſian lixivium, a dirty white precipitate is formed, which becomes yellow in the fire, and becomes white again, on cooling. The Pruſſian lixivium will not precipitate earths from ſolutions of earthy ſalts in water.

f. Liver of ſulphur, or hepatic air, precipitates zinc of a pale coffee colour.

g. Moſt of the precipitates of zinc will be found to be volatile in the fire.

ARSENIC.

Regulus of Arſenic combined with pure Air.

Synonymous Names.

ARSENIC.
- White arſenic.
- Calx of arſenic.
- Flowers of arſenic.
- Lime of arſenic.
- Oxide of arſenic.
- White oxide of arſenic.
- Sublimed white oxide of arſenic.
- Oxygenated arſenic.

Arſenic is found in ſpring waters in the neighbourhood of mines; and as ſuch waters are extremely poiſonous, it is important to detect it. Arſenic is very ſoluble in water, on evaporating the water it cryſtallizes into triangular pyramida cryſtals of a yellowiſh colour.

a. If

a. If it exiſts in any quantity in water, it may be obtained by ſimple evaporation, and either cryſtallizing the arſenic or evaporating to dryneſs, and then expoſing the reſiduum to a red heat, in which it will fly off, giving out a ſtrong ſmell of garlic.

b. On adding a ſolution of liver of ſulphur, or hepatic air, to a ſolution of arſenic in water, yellow clouds will be produced, and a precipitate of the ſame colour will fall to the bottom. The arſenic uniting with the ſulphur, forms orpiment, which is precipitated.

c. A ſolution of blue vitriol being added, clouds of a yellowiſh green colour will be formed, which gradually fall to the bottom. The arſenic, uniting with the copper, forms what is called ſcheel's green.

d. On adding a ſolution of gold to water containing arſenic, the gold, after ſome time, will be precipitated in its metallic ſtate.

e. Upon adding a ſolution of vitriolated iron, a precipitate, of a dirty green colour, will fall to the bottom. The iron combining with the arſenic.

f. Moſt of theſe precipitates alſo will be found volatile in heat, giving the ſmell of garlic as they fly off.

It has been the cuſtom to give mucilaginous drinks to perſons that had ſwallowed arſenic. Milk, fat, oil, butter, &c. But Navier, a Phyſician at Chalones, in France, recommends, from his own experience, the calcarious, or alkaline, livers of ſulphur as the beſt means of counteracting this poiſon. Theſe combine with arſenic in the humid, way ſaturate it, and almoſt diveſt it

it totally of its cauſticity. They act ſtill better, if impregnated with a little iron. He preſcribes a dracm of theſe livers of ſulphur diſſolved in a pint of water, to be drank at various draughts, or five or ſix grains of them to be made into pills; drinking a glaſs of warm water after each doſe: and he recommends the uſe of ſulphureous waters for ſome time after the firſt ſymptoms are removed. He aſſures us, that he has ſeen the happieſt effects of this in removing the tremors and paralytic affections, occaſioned by this poiſon.

SECT. V.

Of the Contents of the Air.

Did not the acid vigor of the mine,
Roll'd from ſo many thund'ring chimneys, tame
The putrid ſteams that overſwarm the ſky;
This cauſtic venom would perhaps corrode
Thoſe tender cells that draw the vital air,
In vain with all their unctuous rills bedew'd *(d)*.

THAT thin, tranſparent, inviſible fluid, called the atmoſphere, which ſurrounds the earth, riſes to a conſiderable height above its ſurface. Although inviſible to us, it is a real ſubſtance, excluding other ſubſtances from the ſpace which it occupies, and, like other fluids, preſſes equally in all directions: its weight is the cauſe of the ſuſpenſion of mercury in the barometer, and is to that of water as 1 to 900, or 15 pounds upon every ſquare inch of the earth.

It chiefly conſiſts of two elaſtic fluids, or gaſes *(e)*, poſſeſſed of very different and oppoſite properties; it contains alſo other gaſes, or ſubſtances, ſuſpended or diſſolved in it, but in ſmall quantity. Theſe two fluids, or gaſes, are diſtin-

(d) Armſtrong's Art of preſerving Health, p. 51.

(e) Gas, from gaſcht (German) an eruption of wind; any matter ſubtilized by heat into an elaſtic aeriform ſtate.

guiſhed by the names of oxygen gas, or the pure part of the atmoſphere, firſt diſcovered by Dr. Prieſtley in the year 1774, and by him called dephlogiſticated air; by others vital air, becauſe it is the only air that is capable of ſuſtaining the vital principle, or of ſupporting the combuſtion of inflammable matter: it is not only the ſole vivifying power, but the parent of acidity, and hence named oxygen *(f)*, from the moſt general property which its baſe poſſeſſes of forming acids, by combining with different ſubſtances.

The other, or noxious portion of the atmoſphere, is termed azote *(g)*, from its known quality of killing animals; hence the noxious part of the atmoſpheric air, which is totally unfit for reſpiration, is called azotic gas, or phlogiſticated air.

The atmoſphere contains other elaſtic fluids analogous to common air, with reſpect to elaſticity and inviſibility; but otherwiſe eſſentially different from each other; ſuch as fixed air, or carbonic acid gas; inflammable air, or hydrogen gas; nitrous gas: beſides a variety of other ſubſtances, reſulting from various exhalations conſtantly emitted from vegetable, animal, and foſſil bodies *(h)*; but it is to the gaſes above enumerated that the following obſervations will be confined.

Atmospherical

(f) Oxygen, οξυς, acidum, or acid, and γινομαι, gignor, to produce, or generate acids.

(g) Azotum, αζωτον, from a privation of, and ζων, life.

(h) Fluoric acid gas, which is diſengaged from native fluate of lime, or vitreous ſpar, by ſulphuric acid. *Muriatic acid gas*, or muriatic acid purified from water, and melted by caloric into an elaſtic

Atmospherical Air, or common Air.

As the two gaſes of oxygen and azote exiſt in the air, merely in a ſtate of intimate mixture, and not in chemical combination, they may be eaſily ſeparated by a ſubſtance poſſeſſing an electric attraction for either of them: thus, if pure mercury be expoſed for ſome time in cloſed veſſels filled with atmoſpheric air, to a heat nearly approaching to ebullition, a quantity of air will diſappear, and the mercury be converted into a red powder, oxyd, or calx of mercury, which will have gained an addition of weight equal to that of the air abſorbed, the mercury having united with the oxygen gas, which leaves the azote. If a bell glaſs be inverted over a ſolution of liver of ſulphur, the air in it will ſuffer a diminution; the oxygen will be abſorbed, and the unabſorbed part will be azote, ſhowing that there are 27 parts of oxygen, and 73 parts of azote in 100 parts of atmoſpheric air.

If the azote in liver of ſulphur be expelled, and the oxygen in the red calx of mercury, or

elaſtic fluid. *Oxygenated muriatic acid gas*, or *dephlogiſticated marine acid*, which is diſengaged during the reciprocal action of native oxyd of manganeſe and muriatic acid, being produced by the tranſition of oxygen from the manganeſe into the muriatic acid. *Ammoniac gas, alcaline air*, or *volatile alcaline gas*, is diſengaged by heat from liquid ammoniac; or, from a mixture of ammoniacal muriate, or common ſal ammoniac, with quick lime. *Hepatic gas*, or ſolphurated hydrogenous gas, is obtained from ſolid alcaline ſulphures, or livers of ſulphur, by decompoſing them with acids. *Phoſphorated hydrogenous gas*, or *phoſphoric gas*, obtained by boiling a lixivium of cauſtic pot-aſh with half its weight of phoſphorus, and receiving the elaſtic fluid diſengaged into glaſſes containing mercury.

precipitate per ſe, and theſe be mixed, atmoſpheric or vital air will be formed.

Oxygen gas conſtitutes about one fourth of the atmoſphere, and is compoſed of light, caloric *(i)*, and oxygen. It is procured, by filling a receiver with the green leaves of vegetables, and inverting it in ſpring water, and expoſing it to the direct rays of the ſun; the leaves yield a conſiderable quantity of oxygen air, which aſcends to the upper part of the receiver, and may be eaſily removed from it for uſe. It is produced from various other ſubſtances, and particularly from nitre, or the metallic calces. One ounce of nitre, expoſed to a red, or rather white heat, in an earthen retort for about four or five hours, will give about 700 cubic inches of oxygen air; as this gas contains a quantity of nitrous acid in the form of vapour, it may be ſeparated from it by agitating the air in lime water.

But it is from manganeſe that the greateſt quantity of this gas may be procured, and it is now the ſubſtance generally uſed. One ounce of good manganeſe will, in a red heat, yield more than two pints and a half, or about 80 cubic inches of elaſtic fluid, one tenth of which is carbonic acid, and the reſt oxygen gas.

(i) Caloric, latent heat, fixed heat, or principle of heat, is only known by its effects; inviſible, imponderable, ſubtile, pervading all bodies, inſinuating itſelf between their particles. It melts ſolid bodies, rarefies fluids, and renders them inviſible. There are but three ſpecies of the undecompounded bodies, which are known to be rendered into the ſtate of gas by coloric; namely, oxygen, hydrogen, and nitragen.

AZOTIC GAS,

Synonymous Names.
- Vitiated air.
- Impure air.
- Phlogiſticated air.
- Phlogiſticated gas.
- Atmoſpheric mephitis.

Is compoſed of caloric, or heat, and a particular baſe, capable of becoming ſolid, called azote. This ſubſtance, united to different baſes, forms alkalies, and may be conſidered as a real alkalizen, or alkalizing principle, in oppoſition to oxygen, which is the principle of acidity. The atmoſphere, therefore, is an immenſe reſervoir of the principles of acidity and alkaleſcency, without being itſelf either acid or alkali.

Azotic gas may be obtained by expoſing certain ſubſtances to atmoſpheric air, which abſorb its oxygen; as ammonia, or the volatile alkali, which is compoſed of azote, united with hydrogen.

CARBONIC ACID GAS,

Synonymous Names.
- Fixed air.
- Solid air of Hali.
- Cretaceous acid gas.
- Mephitic gas.
- Aerial acid.

Is the heavieſt of the aerial fluids, being compoſed of carbon or charcoal, and vital air or oxygen, forming carbonic acid. It is combined with calcareous

careous earth or lime, to which it has a greater affinity than to any other; it conſtitutes nearly half the weight of chalk, limeſtone, or carbonate of lime, &c. converting them into ſaline ſubſtances, or carbonates; it is produced by burning limeſtone, in which the carbonic acid combines with caloric, and flies off in the form of gas, leaving the calcareous earth pure, or in a ſtate of quicklime; it is produced plentifully by fermenting liquors, and is often found in coal mines, called the chalk, or chalk damp: it is formed in caverns, as in the Grotto del Cano, which extinguiſhes flame and deſtroys life. Charcoal, on burning, unites with the oxygen of the atmoſphere, and forms carbonic acid, which is alſo deſtructive to life, and extinguiſhes the light of a candle.

Water abſorbs more than its bulk of this gas, and acquires an acid taſte and ſparkling appearance, and moſt mineral waters owe their ſparkling ſpirituous taſte to this gas; and cyder, beer, and other fermented liquors owe their brightneſs to the carbonic acid which they contain. Water thus impregnated is capable of diſſolving a ſmall quantity of iron, forming a chalybeate water like Spa, or Pyrmont. When fixed alkali is previouſly diſſolved in water, it will then abſorb a greater quantity of carbonic acid than common water, and forms the aqua mephitica alcalina. Limeſtone, or carbonat of lime, cannot be diſſolved but in a very ſmall quantity in water; but pure, or quicklime, can be diſſolved in a conſiderably greater proportion, forming lime-water, which is a teſt of carbonic acid; for it is certainly precipitated

precipitated by that acid in the form of carbonate of lime, or limeſtone. If the water be ſaturated with carbonic acid, it will diſſolve carbonate of lime in conſiderable quantity; and if this ſolution be let fall drop by drop on any ſubſtance, or the carbonic acid eſcapes, the carbonate of lime will be depoſited in the incruſted ſubſtance with calcareous earth, and the ſubſtance is ſaid to be petrified.

HYDROGEN(k) GAS,

OR

INFLAMMABLE AIR,

Inflammable Gas, or Phlogiston of Kirwan.

Hydrogen, which is the baſe of water, combined with caloric, forms hydrogen gas, or inflammable air, diſcovered by Mr. Cavendiſh in 1767. It is the lighteſt of elaſtic fluids, being twelve times lighter than common air; hence employed in air-balloons. It inflames by contact of an ignited body, but will burn only in contact with common, or oxygen, air. It is leſs noxious than the carbonic acid: ſuffers no diminution on mixture with nitrous air. It is produced during the diſſolution of animal and vegetable bodies; hence it

(k) Hydrogenium (ὑδρογενιον, from ὑδωρ, water, and γινομαι, to become, or γενναω, to produce) hydrogen, one of the principles of water. The baſe of that elaſtic fluid, which was formerly called inflammable air.

is

is found to come out of ponds, burying-grounds, and other places containing animal and vegetable matter in a ſtate of decay. Found alſo in the earth where inflammable minerals are contained; when, combining with atmoſpheric air, it ſometimes ſuddenly takes fire to the danger of the miners, and by them called fire-damps. It is obtained from moſt kind of bodies; but the greateſt quantity may be extracted from zinc, by means of diluted vitriolic acid; and from red-hot iron, by paſſing the ſteam of boiling water over its ſurface. In like manner charcoal yields inflammable gas, called hydrocarbonate, combined with the carbonic acid gas. It is this aeriform fluid which floats frequently on marſhes, and being ſet on fire by electricity, or other means, gives riſe to the ignis fatuus. Hydrogen gas has the property of diſſolving and ſuſpending a variety of ſubſtances, as iron, charcoal, ſulphur, phoſphorus, &c. hence we have the names of phoſphoric hydrogen gas, or phoſphuret of hydrogen; ſulphuric hydrogen gas, or ſulphuret of hydrogen; with phoſphorus it is called phoſphorated hydrogen gas, giving out a fœtid ſmell, is improper for reſpiration, and takes fire ſpontaneouſly in coming in contact with the air, accompanied with an exploſion; and probably to the diſengagement of gas, the ignis fatuus, playing about burying-grounds and places where animals are putrifying, may be attributed.

Nitrous acid gas is diſengaged from nitric acid by various combuſtible bodies, eſpecially metals, oils, mucillages, and alcohol. By the action of the nitric acid on theſe ſubſtances it becomes

comes nitrous gas; it extinguiſhes light and deſtroys animals, being totally unfit for animal reſpiration and combuſtion. It is not diſſolved in water, nor does it indicate the leaſt property of an acid: by combination with vital air it affords nitric acid, being itſelf nothing but nitric acid deprived of a part of its oxygen, and conſequently a compound of azote, or nitrogen and oxygen, containing, however, more azote and leſs oxygen than the nitric acid; hence are produced the varieties of this gas, according to the different proportions of azote and vital air. In nitrous gas the azote and oxygen are deprived of all that quantity of caloric and light which they poſſeſſed in the atmoſphere: the oxygen, however, ſtill retains enough of both theſe principles to occaſion a combuſtion, with flame, of ſeveral combuſtible bodies when immerſed in it, as a pyrophorus, or a ſmall portion of ſulphuret of alkali; the nitrous gas is then gradually diminiſhed, and at length pure nitrogen gas remains. In this caſe, the oxygen of the nitrous gas combines with the inflammable body, and thus the nitrogen gas is left in a free ſtate; hence, as has been intimated, it is evident that azote, or nitrogen, conſtitutes the baſis of nitric acid, which obtains its acid property from its combination with oxygen. The mixture of one part of nitrous gas, and four parts of oxygen gas, produces red vapours; becauſe the baſes of each gas mutually combine, and form nitrous acid gas; and the ſuperabundant caloric eſcapes in the ſtate of ſenſible heat. If this operation be inſtituted over mercury, the nitric acid recently formed will remain upon its ſurface, in the form

of nitrous acid gas; but if made over water, it is immediately abſorbed by it.

As azotic gas does not combine with nitrous gas, but remains behind when mixed with it, Dr. Prieſtley employed nitrous gas for aſcertaining the proportionate quantity of oxygen gas in atmoſpheric air. To this end equal parts of nitrous gas and atmoſpheric air are mixed over water, and the greater or leſs diminution of volume of gaſes determines the relative quality of the air examined.

To aſcertain the purity of atmoſpheric or other airs, different kinds of eudiometers have been deſcribed by Prieſtley, Magellan, and Abba Fontana. A ſimple method is deſcribed by Dr. Archer, which is, to have a tube, between two and three feet in length, one third of an inch wide, and open at one end, a meaſure for the airs; a ſcale carefully graduated, according to the contents of the meaſure, into tenths and hundredth parts, ſo that one of the hundredth parts will be about one-ſixth of an inch. The operator is to fill the tube with water, and invert it in a trough of the ſame fluid, and ſet it over a hole on the ſhelf of a trough; the tube will remain perfectly full of water, from the preſſure of the atmoſphere upon the ſurface of the water in the trough. Then fill the air meaſure with water alſo, and if it be the air of the place to be tried, it is only neceſſary to pour out the water, and the air will occupy the ſame ſpace in the meaſure as had before been occupied by the water. Then convey the air, contained in the meaſure, under the ſurface of the water in the trough, to the bottom of the tube; by gently turning

turning the hand, the air in the meaſure may be made to aſcend into the tube, within which it will cauſe a portion of the water to deſcend, exactly correſponding to the quantity of air that was contained in the meaſure. Now mark the ſpace which the air occupies, by applying the graduated ſcale to the outſide of the tube; next take the ſame meaſure full of nitrous air, and introduce it alſo in the ſame manner into the tube; gently agitate the tube, and, after waiting ſome little time, apply the ſcale again, and obſerve what diminution has taken place; for if two meaſures of the ſame air had been introduced, they would occupy of courſe twice as much ſpace within the tube as one would have done. But this will by no means be the caſe in the inſtance before us; for a meaſure of common air, and one of nitrous air, will not occupy as much ſpace as two meaſures of common air, or two of nitrous would, but will fall conſiderably ſhort of it; and it is in proportion to this diminution that the purity of the air is aſcertained, the greater diminution denoting the greater purity of the air. Thus, if one meaſure of nitrous air be added to one of any other kind of air, and the diminution be $1\frac{26}{100}$, the meaſure of the teſt will be 1.26. The nitrous acid being compoſed of nitrous air and oxygen air, whenever theſe two airs meet, they put off their aëriform ſtate, and become liquid nitrous air; which, as it mixes with the water, no longer impedes the aſcent of that fluid in the tube.

The ſpecific gravity of different kinds of air:

A cubic inch of atmoſpheric air weighs	385.
dephlogiſticated -	420.
phlogiſticated -	377.
fixed - -	570.
inflammable -	35.
nitrous - -	399.

SECT VI.

Directions for collecting and diſtinguiſhing Foſſil *ſubſtances, including* Salts, Earths, Metals, *and* Inflammables.

TO write particularly upon theſe objects of natural hiſtory is not the preſent deſign, but to give ſuch general inſtructions, as may aſſiſt a traveller, in the choice of foſſil bodies, till fuller information can be procured, which the numerous modern authors on theſe ſubjects afford. In the arrangement adopted, I have principally had in view that by Dr. Babington; and his own and the other references annexed to the deſcriptive hiſtories, will point out the principal writers on the ſubjects of diſcuſſion.

The traveller ſhould be furniſhed with flint and ſteel, and the mineral acids; at leaſt with the nitrous, and volatile acids; a hammer alſo will be neceſſary, to break bodies too bulky to bring away.

A blow-pipe is likewiſe an uſeful article; by blowing long and forcibly through ſuch an inſtrument, upon the flame of a candle, by which the point of the flame may be directed againſt the body to be examined, it will frequently diſcover whether it is a calcareous, vitrifiable, or refractory ſubſtance; and for greater preciſion, experiments

ments for the ſame end may be afterwards made at the fire-ſide, on a charcoal fire. Even a tobacco-pipe may be uſed as a crucible in a common fire.

The collector ſhould alſo attend to as many of the following particulars as poſſible.

1. When any article is collected, mark it by a number, or ſome ſign of diſtinction, referring to a catalogue, with all the particulars that may be known relative to it; as,

2. Where it was found.

3. In what quantity, whether ſcarce or abundant.

4. If on the ſurface of the earth, or at what depth.

5. In what poſition, whether horizontal, perpendicular, &c. And with what other foſſil bodies it was found; as clay, ſtone, ſlate, mineral, &c.

6. Whether in ſtrata, or looſe nodules.

7. The depth and thickneſs of the ſtrata, how they incline, or to what points of the compaſs they tend; or if level or horizontal; whether they have perpendicular or horizontal fiſſures, and what foſſil bodies are contained in theſe fiſſures.

8. All high mountains and hills, eſpecially their ſides, are to be ſearched; the ſhores alſo of the ſea, with their banks, and the cliffs adjacent, and the falls of caſcades, rivers, and great gullies.

9. The ſituation of mines, pits, and quarries, whether in a valley or hill; and the diſpoſition of the ſtrata, whether horizontal, inclining, &c. their thickneſs, and the depth they lie; and what other foſſils are imbedded in the ſtrata, or in the neighbouring caverns, fiſſures, partings, &c.

10. The

10. The waters of mines ſhould be examined, whether pure, taſteleſs, purgative, vitriolic, or chalybeate, &c.

11. The damps and ſteams of mines, and what are the conſequences or effects of them; in what ſeaſons and ſtate of the air they are chiefly obſerved; and what temperature the air bears in different depths of ſuch mines.

12. The account given by the natives, inhabitants, miners, workmen, &c. who may be acquainted with the ſubject.

CLASS

CLASS I.

SALTS.

1. Simple Salts,

Having radicals, which are simple and known.

A. *ACID.* All acids reſemble one another in taſte; in their manner of giving a red colour to vegetable ſubſtances; in their common tendency to combine with earths, alkalis, and metallic oxyds.

a. *Carbonic.*

Synonymous Names.
- Gas ſylveſtre.
- Spiritus ſylveſtris.
- Fixed air.
- Aërial acid.
- Atmoſpheric acid.
- Mephitic acid.
- Cretaceous acid.
- Carbonaceous acid.

See Sect. V. page 45, 85.
Anal. Carbon. 17. Oxygen 83. *Chaptal.*

b. *Boracic.*

Synonymous Names.
- Volatile narcotic ſalt of vitriol.
- Sedative ſalt.
- Acid of borax.
- Boracine acid.

This

This acid is dry, cryſtallized in hexahædral plates, almoſt inſipid, ſcarcely ſoluble, fuſible with ſilices into a glaſs, very feeble in its affinities, and liable to yield up the terrene and alcaline baſes to almoſt any other acid.

	Synonymous Names.
c. *Sulphuric.*	Acid of ſulphur. Vitriolic acid. Oil of vitriol. Spirit of vitriol.
Sulphureous.	Sulphureous acid. Volatile ſulphureous acid. Phlogiſticated vitriolic acid. Spirit of ſulphur.

The *ſulphuric acid* is formed from ſulphur and oxygen, by the combuſtion of the ſulphur; inodorate, twice as heavy as water, and leſs volatile, very cauſtic; affording ſulphureous acid gas and ſulphur, by its decompoſition with *red hot coal*, metals, &c. forming ſulphates *(l)* with the earths, alkalis, and metallic oxydes.

Anal. Sulph. 72. Oxygen 28. *Berthollet.*

Sulphureous acid is highly odorate, volatile, gaſeous, deſtructive to blue vegetable colours, and fit to cleanſe away ſpots made with theſe colours on a white ground; abſorbing, by ſlow degrees, the oxygen from the air, and from various

(l) Salts formed by the combination of the ſulphuric acid with different baſes, decompoſable by charcoal, &c. into ſulphures.

acids and oxydes; forming ſulphites *(m)* with the earthy and alkaline baſes.

2. Compound Salts.

A. *Base POT-ASH*, or *Vegetable, caustic, fixed alkali.* The impure ſalt, containing alkali, which is obtained by evaporating the aqueous ſolution of wood aſhes to dryneſs; it is then ſolid, white, cryſtallized in rhomboidal plates.

a. *Carbonate* of Pot-ash*, or pot-aſh combined with carbonic acid;

Synonymous Names.
- Fixed ſalt of tartar.
- Vegetable fixed alkali.
- Aërated vegetable fixed alkali.
- Cretaceous tartar.
- Mephitic tartar.
- Mephite of Pot-aſh.
- Nitre fixed by itſelf.
- Alkaheſt of Van Helmont.

It forms regular cryſtals, which repreſent tetrahædral priſms, terminating in ſhort tetrahædral points; has an urinous taſte, and changes the ſyrup of violets green.—*Anal.* Acid 23. Alkali 70. Water 5. *Bergman.*

(m) Salts formed by the combination of the ſulphureous acid with different baſes, and yield by the contact of almoſt any acid, the ſmell of burning ſulphur with effervescence.

* A ſalt formed by the union of carbonic acid with different baſes.

b. *Muriate*

b. *Muriate* of Pot-ash,* or Febrifuge ſalt of Sylvius; compoſed of muriatic acid and pot-aſh; is found in its native ſtate in ſea-water; in the mother earth of nitre manufactories; in various ſalt-ſprings; has rhomboidal or octahædral cryſtals.—*Anal.* Acid 31. Alkali 60. Water 8. *Bergman.*

c. *Nitrate† of Pot-ash, Nitre,* or *Saltpetre;* formed by the combination of nitric acid and pot-aſh. It is found in the fiſſures of lime-ſtone kilns, near Molfetta, in the kingdom of Naples; in various waters, and even in rain; and in the freſh juices of many plants.—*Anal.* Acid 33. Alkali 49. Water 18. *Bergman.*

d. *Sulphate‡ of Pot-ash.*

Synonymous Names.
- Vitriol of pot-aſh.
- Sal de duolus.
- Vitriolated tartar.
- Arcanum duplicatum.
- Sal polychreſt of Glaſer.

compoſed of the ſulphuric acid and pot-aſh: ſcarcely ever found native.—*Anal.* Acid 40. Alkali 52. Water 8. *Bergman.*

* Salts formed by the union of the muriatic acid with different baſes.

† Salts formed by the combination of the nitric acid with different baſes.

‡ Salts formed by the combination of the ſulphuric acid with different baſes.

B. *Base SODA.* Cauſtic ſoda, marine alkali, or mineral alkali united to carbonic acid, which is obtained from the aſhes of the kali ſpinoſum, and of many ſea plants.

Synonymous Names.

a. *Carbonate of Soda,*
- Natrum or natron.
- Baſe of marine ſalt.
- Marine or mineral alkali.
- Cryſtals of ſoda.
- Cretaceous ſoda.
- Aërated ſoda.
- Efferveſcent ſoda.
- Mephite of ſoda.
- Aërated mineral fixed alkali.
- Efferveſcent mineral fixed alkali.
- Chalk of ſoda.

is a perfect neutral ſalt, conſiſting of the carbonic acid, and pure or cauſtic ſoda; has rhomboidal cryſtals. Found in Hungary, Egypt, Perſia, the Eaſt Indies, and China; is found ready formed on the ſurface of the earth.—*Anal.* Acid 16. Alkali 20. Water 47. *Delamethrie.*

b. *Borate* of Soda.* Greyiſh and greeniſh white, moſtly in tabular, ſix-ſided priſms. Found on the ſnow mountains of Tibet, &c.—*Anal.* Acid. 34. Alkali 17. Water 47. *Delamethrie.*

c. *Muriate of Soda,* or marine ſalt, or culinary ſalt. Cryſtallized in cubes, amorphous,

* Borax Tinkal.

fibrous

fibrous, compact. *Anal.* Acid 52. Alkali 42. Water 6. *Bergman.*

d. *Sulphate of Soda.* Glauber ſalt, or vitriol of ſoda. Sulphuric acid with ſoda.—*Anal.* Acid 27. Alkali 15. Water 58. *Bergman.*

c. *Nitrate of Soda.* Cubic nitre, or rhomboidal nitre formed by the combination of the nitric acid and ſoda.

Synonymous Names.

C. *Base*, AMMONIAC,
- Cauſtic volatile alkali.
- Fluor volatile alkali.
- Volatile ſpirit of ſal ammoniac.

a. *Carbonate of Ammoniac*,
- Ammoniacal chalk.
- Cretaceous ammoniacal ſalt.
- Concrete volatile alkali.
- Ammoniacal mephite.
- Engliſh ſal volatile.

Sal ammonia united to carbonic acid.—*Anal.* Acid 45. Alkali 43. Water 12. *Bergman.*

b. *Muriate of Ammoniac*, ſalmiac, ſal ammoniac, or muriated ammoniac, compoſed of the muriatic acid and ammoniac, or the volatile alkali. It is found ready formed near the craters of volcanos, and in Tartary. In Egypt it is prepared in great abundance by ſubliming the ſoot, obtained from the combuſtion of animal excrements.—*Anal.* Acid 52. Alkali 40. Water 8. *Delamethrie.*

c *Nitrate*

c. *Nitrate of Ammoniac*; ammoniacal nitrous ſalt, or ammoniacal nitre; a combination of the nitric acid and ammoniac. It is generated ſpontaneouſly in the mother earth, or ſaltpetre manufactories, though but in ſmall quantities; mixed with nitrate of pot-aſh.—*Anal.* Acid 46. Alkali 40. Water 14. *Delamethrie.*

d. *Sulphate of Ammoniac*; ammoniacal vitriolic ſalt, ammoniacal ſalt (ſecret of Glauber's) or ammoniacal vitriol, formed by the ſulphuric acid and ammoniac; concrete.—*Anal.* Acid 42. Alkali 40. Water 18. *Delamethrie.*

CLASS

CLASS II.

EARTHS.

EARTHS are ſuch bodies as are inodorous, inſipid, pulverulent or friable, not flammable, ſcarcely ſoluble in water, unchangeable in the fire; when heated, either by themſelves or with inflammable bodies, do not acquire metallic ſplendour.

I. Homogeneous or simple Earths.

Thoſe earths which have not hitherto been decompounded, are termed elementary, homogeneous or ſimple earths. Of this kind are the following nine. A. Lime or calcareous earth. B. Stranthian earth. C. Baryte or ponderous earth. D. Magneſia or earth of bitter ſalt. E. Alumine or argillaceous earth. F. Silex or ſiliceous earth. G. Adamantine earth. H. Jargon earth. I. Sidnean earth; this laſt, however, is doubtful as an elementary earth.

A. *LIME.*

a. *Pure lime*, cauſtic or pure calcareous earth, in a ſtate of mixture with aërated calcareous earth, has hitherto been found only at Bath, and near the craters of ſome ancient volcanos.

b. *Carbonate*

Synonymous Names.

b. *Carbonate of Lime,*
- Chalk.
- Limeſtone.
- Calcareous mephite.
- Aërated calcareous earth.
- Effervescent calcareous earth.
- Calcareous ſpar.
- Cream of lime.
- Calcareous carbonate.

effervesces with acids, convertible into pure, or quick-lime, by calcination; is then ſoluble in 680 times it's weight of water, forming lime-water, in which action it gives out it's latent heat, cauſing the maſs to become hot.—*Anal.* Carbonic Acid 34. Lime 55. Water 11. *Bergman.* Here may be included ſtalactites, marble, limeſtone, chalk, zuſa, ganil.

c. *Swine-ſtone,* carbonated lime impregnated with petroleum; colour, Iſabella yellow, bluiſh, or blackiſh, or yellowiſh grey, is opaque, compact, and partly of a glittering appearance; when broken, exhibits a ſplintery or conchoidal appearance; when pounded or ſcraped, it emits an urinous or alliaceous ſmell. *Anal.* Carbonate of lime impregnated by petroleum. *Kirwan.*

d. *Sidero-Calcite,* pearl ſpar: colour of a white greyiſh, yellowiſh, or reddiſh white; found amorphous, cellular, rounded, but moſtly cryſtallized; externally metallic or pearly; effervesces with acids. *Anal.* Carbonated lime 60. Oxyde of magneſia, 35. Iron 5. *Woulfe.*

e. *Baryto-*

e. *Baryto-calcite*, of a dark, or light grey, almoſt white; rounded form, or cryſtallized in quadrangular priſms; effervesces with acids. *Anal.* Carbonated lime 92. Carbonated baryte 8. *Bergman.*

f. *Muri-calcite*, Compound ſpar, of an olive colour, clayey appearance; ſtony, amorphous,* and cryſtallized. *Anal.* Carbonated lime 52. Carbonated magneſia 45. Iron and manganeſe 3. *Klaproth.*

g. *Argentine*, Scheifer ſpar, amorphous, foliated. *Anal.* Carbonated lime, with magneſia, argill, and iron. *Kirwan.*

h. *Supersaturated with carbonic acid*, Dolomite, or elaſtic marble, pure or greyiſh white, yellowiſh white, or light red; ſurface rough and uneven; fracture conchoidal or granularly foliated. Effervesces ſlowly with acids; phoſphoreſces when ſtruck in the dark, or when laid on red-hot iron. *Anal.* Lime 44, 29. Carbonic acid 46, 1. Argill 5, 86. Magneſia 1, 4. Iron 0, 074. *Saussure, jun.*

Synonymous Names.

i. *Fluate*† *of lime*, { Fluor ſpar. / Vitreous ſpar. / Cubic ſpar. / Phoſphoric ſpar. / Sparry fluor.

Nearly inſoluble in water, does not effervesce with acids, nor ſtrike fire with ſteel; is more or leſs tranſparent, and exhibits various colours; it's

* Amorphous, ex α priv. et μορφη, forma.

† A ſalt formed by the combination of the fluoric acid with different baſes.

ſurface, when ſcratched with a knife, appears greaſy; it generally crackles and phoſphorates by heat, except the colourleſs, which becomes electric when rubbed; and the coloured kinds loſe their colour by heat; but by a ſtrong heat is melted. *Anal.* Fluor acid 16. Lime 57. Water 27. *Scheele.* Oxyde of manganeſe 3. Oxyde of iron 1. Pot-aſh 18. *Vauquelin.*

Synonymous Names.

k. *Phosphate of lime.* { Earth of bones. / Calcareous phoſphate. / Animal earth. }

Apatite, colour various, olive green, bluiſh, yellowiſh and chocolate brown; found cryſtallized in three and ſix-ſided priſms; in ſix and eight-ſided plates, variouſly modified, ſhining and of a greaſy luſtre, texture laminated, ſemi-tranſparent. *Anal.* Phoſphoric acid; lime 55. *Klaproth.*

Synonymous Names.

l. *Sulphate of lime,* { Vitriol of lime. / Calcareous vitriol. / Selenite. / Gypſum. / Plaſter of Paris. }

does not effervеſce with acids; found in moſt parts of the earth and in water; ſoluble in about 500 times it's weight of water; it reſembles the calcareous ſpar, and is found in ſolid maſſes, of particular form, cryſtallized into hexagonal* priſms, terminating in an edge, or in rhomboids, or cuneform, tabular or lenticular; texture fi-

* Hexagonal, ex ἕξ, ſex; et γωνία, angulus.

brous

brous, or laminated. *Anal.* Sulphuric acid 16. Lime 57. Water 27. *Scheele.*

B. *STRONTIAN*, or Scottiſh earth, ſo called from Strontian in Scotland; never found pure. Sulphate of Strontian is ſaid to have been found in great quantities in the neighbourhood of Briſtol. At Ham-green alſo, a variety of this rare production is found breaking through the ſoil, in ſuch large maſſes, that it has been made uſe of in mending the roads.

a. *Carbonate of Strontian, strontianite*; hitherto found in a mild ſtate united to fixed air; in ſolid maſſes, of a fibrous texture, ſtriated. Like limeſtone, it loſes it's air in a ſtrong heat, and then forms a lime, heavier than common lime, and more ſoluble in water. This lime-water has the ſingular property of affording, when ſaturated, compreſſed rhomboidal cryſtals, which are nothing but pure lime. *Anal.* Carbonic acid 26, 5. Stronthian 73,5. *Kirwan.*

C. *BARYTE*, Ponderous earth; never found in a ſeparate or pure ſtate, but with the ſulphuric acid, in the ſtate of ponderous ſpar or ſulphate of baryte; or with the carbonic acid, carbonate of baryte, or mixed with other earths.

Synonymous Names.

a *Carbonate of Baryte*,
- Barotic or ponderous chalk.
- Aërated ponderous earth.
- Effervescent barotes.
- Barotic mephite.
- Barolite.

found in ſolid maſſes, and cryſtallized, hexagonal priſms, terminating in ſix-ſided pyramids; texture ſhining, radiated, fibrous; brittle and ſemi-tranſparent; effervefces with diluted nitric and muriatic acids, and is entirely diſſolved by them. Expoſed to a ſtrong heat, it parts with it's carbonic acid, and combines with heat, and like lime is ſoluble in water. *Anal.* Acid 20,8. Baryte 78,6. *Withering.*

Synonymous Names.

b. *Sulphate of Baryte,* { Heavy or ponderous ſpar. Barotic vitriol. Baroſelite.

found in ſolid maſſes, cryſtallized, approaching to an earthy ſtate, and the tranſparent kind is electric; is four times heavier than water. When expoſed to heat, and to the ſun, it has the property of abſorbing light, which it emits in the dark. Like metals, it is precipitated from it's combinations with the different acids, by the pruſſiate of pot-aſh*. *Anal.* Sulphuric acid 32,8. Baryte 67,2. *Withering.*

Bolognian Stone. Anal. Sulphate of baryte 62. Silex 16. Argill 15. Sulphate of lime 6. Water 2. *Arvidson.*

c. *Liver-ſtone,* of a greyiſh, or greyiſh black colour; texture foliated, ſtriated; does not effervefce with acids; when heated, or rubbed, it

* Salts formed by the union of the Pruſſic acid, or colouring matter of Pruſſian blue, with different baſes.

emits

emits a ſmell of liver of ſulphur. *Anal.* Sulphate of baryte 38. Silex 33. Sulphate of argill 22, Sulphate of lime 7. Petroleum 3. *Bergman.*

D. *MAGNESIA* is never found naturally pure, but generally combined with the ſulphuric acid, in the form of Epſom ſalt, ſal amarus, or ſulphate of magneſia.

a. *Calci-murite*, amorphous, earthy; colour blue or olive green; of the conſiſtence of clay. *Anal.* Silex 50. With carbonate of magneſia and iron.

b. *Argillo-murite*, of a looſe earthy texture; greeniſh yellow colour, conſiſtence earthy. *Anal.* Magneſia, 13. Silex 50. Argill 10. Lime 3. Oxyde of iron 0,9. Water 13. *Kirwan.*

c. *Silici-murite*, Martial muriatic ſpar, amorphous, foliated. *Anal.* Silex 50. With carbonate of magneſia and iron. *Kirwan.*

d. *Talc*, Talcite, Venetian talc, or Shiſtoſe talc. Generally of a greeniſh or ſilvery white luſtre, like mother of pearl, ſoft and ſemi-tranſparent; texture laminated, foliated, and greaſy, or of a ſoapy feel to the touch; does not effervesce with acids. *Anal.* Magneſia 44. Silex 50. Argill 6.

e. *Lapis Ollaris*, Pot-ſtone, or Indurated talc; of a pale yellowiſh and greeniſh grey, reddiſh grey, or white colour, with micaceous particles; found in

in amorphous maſſes, of a lamellated texture; feels greaſy, does not adhere to the tongue; becomes hard in the fire. *Anal.* Magneſia 38. Silex 38. Argill 7. Iron 5. Carbonate of lime 1. And a trace of fluoric acid. *Weigleb.*

f. *Steatites*, Fatty ſtone, of a greyiſh, yellowiſh, or greeniſh white, with a ſhade of green colour; found in amorphous maſſes, of an earthy, ſplintery texture, lamellated; the thin lamina ſemi-tranſparent; ſoapy to the touch, does not adhere to the tongue. *Anal.* Magneſia, 20,84. Silex 48,42. Argill 14. Iron 1. Air and Water 16. *Klaproth.*

g. *Serpentine*, of a deep, blackiſh, olive green, ſeldom yellow, ſometimes crimſon red, bluiſh and greeniſh grey; generally exhibits various colours, like the ſkin of a ſerpent; found in ſolid maſſes; when broken has a dull appearance, ſmooth to the touch, and takes a poliſh. *Anal.* Magneſia 35. Silex 45. Magnetic iron 14. Carbonate of lime 6,25. Argill 0,25. *Knock.*

h. *Chlorite*, of a light and dark greeniſh colour, or greeniſh brown; of a ſcaly texture and glittering appearance; feels greaſy or earthy; gives an earthy ſmell, when breathed on. It melts into a dark, black, compact ſlag, and thus becomes magnetic. *Anal.* Magneſia 0,39,47. Silex 0,415. Argill 0,0613. Lime 0,015. Iron 0,1015. Air and water, 0,015. *Hoepfner.*

i. *Asbestos*,

i. *Asbestos*, of a leek or olive green colour, or grey; feels greasy; breaks into parallel, striated, and sometimes curved, fibres or splinters; does not effervesce with acids. *Anal.* Carbonate of magnesia 16. Silex 63,9. Carbonate of lime 12,8. Argill 1,1. Oxyde of iron 6. *Bergman.*

k. *Amianthus*, of a green or greyish, or silvery, white, sometimes olive green; breaks into flexible, and mostly parallel, filaments; soft and greasy, somewhat transparent. *Anal.* Carbonate of magnesia 18,6. Silex 64. Carbonate of lime 6,9. Barosselenite 6. Argill 3,3. Oxyde of iron 12. *Bergman.*

l. *Suber Montanum*, cork-like, or resembling brown wood; it is opake, elastic, of a flaty curve, and irregularly fibrous texture. *Anal.* Carbonate of magnesia 22. Silex 62. Argill 2,8. Carbonate of lime 10. Oxyde of iron 3,2. *Bergman.*

m. *Anctynolite*, amorphous, lamellated, striated, and fibrous; in compressed hexahedral prisms with smooth surfaces. *Anal.* Carbonate of magnesia 20. Silex 64. Carbonate of lime 9,3. Argill 2,7. Oxyde of iron 4. *Bergman.*

n. *Jade*, of a dark leek green or bluish colour; surface smooth, sometimes uneven; in detached round masses, or inhering in rocks; of a splintery texture; feels greasy. *Anal.* Carbonate of magnesia

magnesia 38. Silex 47. Carbonate of lime 2. Argill 4. Oxyde of iron 9. *Hoepfner.*

o. *Baikalite*, crystallized in tetrahedral prisms, entire or truncated, with oblique prisms; and in hexahedral prisms. *Anal.* Magnesia 30. Silex 44. Lime 20. Oxyde of iron 6. *Lowitz.*

p. *Boracite*, found only in the mountain of Kalkberg, near Luneburg; colour greyish white, inclining to purplish; crystallized in cubes, with truncated edges and angles; does not effervesce, or dissolve in acids, unless heat be applied. *Anal.* Boracic acid 68. Magnesia 13. Lime 11. Silex 1. Argill 1. Iron 1. *Westrumb.*

E. *ARGILL*, or Alumine, aluminous, or argillaceous earth; the earth contained in sulphate of alumine, or common alum, has no taste, and is insoluble, hardens in the fire, although divisible in water, contracting in it's dimensions, and when exposed to an extreme heat, becomes so hard as to strike fire with steel.

a. *Sulphate of Alumine*, or alum, is an earthy salt, which consists of the sulphuric acid and alumine.

b. *Carbonate of Argill*, Lac lunæ. Colour pure white, found in nests, in compacted, rounded, kidney-form masses. Consists of carbonic acid, argill and some lime. *Schreiber.*

c. *Clay*, *Pipe Clay*, white, very little greasy to the

the touch, friable and ſtains the fingers ſlightly, diffuſible in water and ductile; effervesces a little with acids. *Anal.* Carbonate of argill and ſilex.

d. *Porcelain Clay*, white, and greyiſh, ſoft to the touch; when mixed with and moulded into veſſels, and expoſed to a ſtrong heat, aſſumes the appearance of a ſemi-tranſparent glaſs; ſtrikes fire with ſteel; is not acted upon by acids; and preſerves it's ſhape. *Anal.* Argill 60. Silex 20. *Wedgewood.*

e. *Lithomarga*, Mountain ſoap, friable, indurated, amorphous; texture frequently poliſhed, ſlaty, conchoidal; colour white, yellowiſh or reddiſh white; emits a phoſphoreſcent light, when rubbed in the dark with a pin; ſoft and ſoapy to the touch, abſorbs water rapidly, and retains it for a long time. *Anal.* Argill 11. Silex 60. Carbonate of lime 5,7. Carbonate of magneſia 0,5. Oxyde of iron 4,7. Air and water 18. *Bergman.*

f. *Fullers Earth*, greeniſh white colour, or greyiſh white, receives a poliſh from friction; does not adhere to the tongue; feels ſomewhat greaſy; falls into powder in water. *Anal.* Argill 0,25. Silex 0,51. Carbonate of lime 0,03. Carbonate of magneſia 0,007. Oxyde of iron 0,03. Moiſture and air 0,15. *Bergman.*

g. *Bole*, particles very ſmall, and cohere

ſlightly;

ſlightly; ſoftens eaſily in water, but is not fit for pottery; feels greaſy; acquires a poliſh by light friction; of different colours, the red called *Armenian bole*, uſed for red pencils. The *Terra Lemnia*, which belongs to this tribe, contains ſilex, argill, magneſia, calcareous earth, iron and water. The indurated bole, or *red chalk*, is hard and brittle, and intimately united with oxyde of iron. *Anal.* Argill 19. Silex 47. Carbonate of lime 5,4. Carbonate of magneſia 6. Oxyde of iron 5,4. Water and air 17. *Bergman.*

h. *Tripoli*, ſo called from the place whence it was firſt brought to us; colour whitiſh, grey or ochre yellow; is found ſolid; has a dull and earthy appearance when broken; is ſoft and ſandy between the teeth; abſorbs water with a noiſe, during which air bubbles are expelled; does not effervеſce with acids, unleſs mixed with marl, which it often is. *Anal.* Argill 7. Silex 90. Iron 3. *Haſſe.*

i. *Lepidolite*, amorphous, foliated. *Anal.* Argill 38,25. Silex 54,5. Oxyde of iron and manganeſe 0,075. Water and air 2,5. *Klapr.* Silex 54. Alumine 20. Fluate of lime 4. Oxyde of manganeſe 3. Oxyde of iron 1. Potaſh 18. *Vauquelin. Lelievre.*

k. *Sappare*, Cyanite, compoſed of long laminæ; white, with ſhades of ſky or Pruſſian blue; luſtre like mother of pearl; it is rather brittle, ſemitranſparent, ſoft, and can be ſcraped with a knife; feels to the touch like talc. *Anal.* Argill 67.

Silex

Silex 13. Magnefia 13. Iron 5. Lime 2. *Saussure, jun.*

l. *Mica*, lamellated ſtructure, the laminæ ſometimes of conſiderable ſize, ſemi-tranſparent, and the *Muscovy glass* in theſe laminæ is quite tranſparent, and is uſed for windows and lanterns; flexible and elaſtic; is found of various colours, from ſilver white to blackiſh. *Anal.* Argill 28. Silex 38. Magnefia 20. Oxyde of iron 0. *Kirwan.*

m. *Micarelle*, found in granite; of a browniſh black colour; texture foliated. *Anal.* Argill 63. Silex 29. Oxyde of iron 7. *Klapr. Anal.* Argill 63. Silex 29. Oxyde of iron 7. *Kirwan.*

n. *Hornblende*, Baſaltic hornblende, Common hornblende, Schiſtoſe hornblende, found of various degrees of hardneſs, but never ſo hard as to ſtrike fire with ſteel; blackiſh or deep green colour; cryſtallized in ſix-ſided priſms, the ends acuminated by three faces; and alſo terminating variouſly; cryſtals internally ſhining; the texture exhibits diverging radiations. *Anal* Argill 27. Silex 58. Lime 4. Magnefia 1. Lime 9. *Bergman.* Conf. *Kirwan.*

o. *Resplendent Hornblende*, Labrador hornblende, Schiller ſpar, colour greyiſh black, ſometimes with a ſhade of copper red, reſembling Labrador ſtone; texture lamellated; the lamellæ curved. *Anal.* Argill 17. Silex 42. Magnefia 11. Iron 28. *Gmelin.*

p. *Basalt*, Trap, Wakken, Mullen, Krag; colour blackiſh green, and ſometimes deep black; cryſtallized in ſix and eight ſided priſms, cuneated on the ends; it's ſurface ſhining on being broken; when longitudinally, the lamellæ are parallel. *Anal.* Argill 15. Silex 50. Carbonate of lime 8. Iron 28. Magneſia 2. *Bergman.* Conf. *Withering.*

q. *Calp*, ſlaty, amorphous. *Anal.* Argill, ſilex, and iron, with fifty per hundred weight of carbonate of lime. *Kirwan.*

r. *Argillaceous Schistus*, Argillite, Killas, Grapholite; colour bluiſh or bluiſh grey, purpliſh or black; amorphous and ſlaty. *Anal.* Argill 25. Silex 60. Magneſia 9. Iron 6. And ſome petroleum. *Kirwan.*

ſ. *Novaculite*, Turkey hone, amorphous and ſlaty.

F. *SILEX*, is rough to the touch, cuts glaſs, and ſcratches metals; is infuſible, and inſoluble in moſt acids, except the fluoric, forming Derbyſhire ſpar; ſoluble in alkalies, with the aſſiſtance of heat forming glaſs; found in moſt ſtones that ſtrike fire with ſteel.

a. *Diamond*, cryſtallized in octohedrons* and their varieties; and in dodecahedrons† and icoſa-

* Octohedrons, ex ὀκτω, eight; et ἑδρα, baſis.

† Dodecahedral, ex δωδεκα, twelve; et ἑδρα, baſis.

hedrons

hedrons* and their varieties, indeterminate; found in a ſandy earth in the hither peninſula of India, in the iſland of Borneo, and in the Brazils. Diamonds are of a lamellated texture, and may be ſplit by a blow in a proper direction; cut all other ſubſtances, and take a moſt exquiſite poliſh.

b. *Sapphire* is next in hardneſs to the diamond; the ſtone worn by the high prieſt of the Jews; of a ſky blue colour, ſometimes inclining to pink, ſometimes almoſt white, and then called *lux sapphire*; it is found cryſtallized in lengthened hexahedral priſms joined baſe to baſe. *Anal.* Silex 35. Argill 58. Carbonate of lime 5. Iron 2. *Bergman.*

c. *Topaz of Brazil*, colour various, the true oriental is almoſt colourleſs, or pale yellow; the Brazil is of a fine, yellow, tranſparent amber, cryſtallized in tetrahedral rhomboidal priſms, terminated by tetrahedral pyramids.

d. *Topaz of Saxony*, cryſtallized in four-ſided rhomboidal priſms, terminating from the ſides in flat four-ſided pyramids, longitudinally ſtriated, and of a foliaceous texture. *Anal.* Silex 52. Argill 44. Lime 2. Iron 0,03. *Wiegleb.*

e. *Beryl of Siberia*, colour greyiſh green, verging on the apple green, more rarely bluiſh green; cryſtallized in equi-angular hexahedral priſms longitudinally ſtreaked: becomes electric by

* Icoſahedrons, ex εικοσι, viginti; et εδρα, baſis.

friction,

friction, but one of it's poles is attractive whilſt the other is repulſive. *Anal.* Silex 61. Argill 29. Calx 2. Iron $\frac{1}{2}$. Seven parts loſt. *Herman.*

f. *Ruby*, of a crimſon, deep red colour, which it retains in the fire; cryſtallized in octohedrons and their varieties, and double four-ſided, ſeldom ſix-ſided pyramids. *Anal.* Silex 16. Argill 76. Lime 1. Iron 3. *Klaproth.*

g. *Emerald*, Smaragdite, of a pure green colour, which it loſes in the fire, as well as it's weight; cryſtallized in hexagonal priſms, either perfect or truncated on the angles or edges, terminating in truncated pyramids; the cryſtals moſtly ſmooth on the ſurface, ſhining and tranſparent in various degrees. The light green coloured oriental emerald is reckoned the beſt. *Anal.* Silex 24. Argill 60. Lime 8. Iron 6. *Klaproth.* The Smaragdite has been analyſed by *Vauquelin.* The *Green and White Smaragdite* of Corſica. Silex 50. Alumine 11. Magneſia 6. Lime 13. Oxyde of iron 5,5. Oxyde of copper 1,1. Oxyde of Chrome 4. The *green* Smaragdite. Silex 51. Alumin 13,5. Magneſia 5. Lime 14,5. Oxyde of iron 8. Oxyde of copper 0,5. Oxyde of Chrome 4. The *Grey.* Silex 50. Alumine 7. Magneſia 8. Lime 17. Oxyde of iron 14,5.

h. *Aqua Marine*, Aigue marine, cryſtallized in large, perfect, ſix-ſided priſms; colour of a pale green *Anal.* Silex 64. Argill 24. Lime 8. Iron 1,5. *Bindheim.*

i. *Crysolite*,

i. *Crysolite*, colour yellowiſh green, inclining to yellowiſh brown; cryſtallized in hexahedral priſms with correſponding pyramids. *Anal.* Silex 15. Argill 64. Lime 17. Iron 1. *Achard.*

k. *Hyacinth*, of a peculiar yellowiſh red; in general retain their colour in fire; cryſtallized in dodecahedrons with unequal rhombic faces. *Anal.* Silex 25. Argill 40. Carbonate of lime 20. Iron 13. *Bergman.*

l. *Hyacinth of Vesuvius*, is generally of a deeper colour than the oriental, and is leſs hard and more fuſible. Cryſtallized in tetrahedral priſms, truncated at their angles, terminated by tetrahedral pyramids, truncated at their ſummits.

m. *Olivin*, found imbodied in baſaltes; of an olive green colour; cryſtallized in ſix, ſeldom in four, ſided priſms; moſtly rectangular, with or without pyramids. The cryſtals are longitudinally ſtriated; have a vitreous appearance. *Anal.* Silex 54. Argill 40. Iron 4. *Gmelin.*

n. *Garnet*, Carbuncle, of a red colour, or ſome of it's ſhades, a greeniſh white, and of various degrees of tranſparency; does not loſe it's colour in the fire, but becomes heavier, and is ſtrongly attracted by the magnet. Cryſtallized and preſenting either 12 rhomboidal planes, or 24 trapezoidal, or 36 planes, of which 12 are rhombs, and the 24 others elongated hexagons, interpoſed between theſe rhombs, and ſome other variations. *Anal.* Silex 48. Argill 30. Lime 11. Iron 10. *Achard.*

o. *White*

o. *White Garnet*, Veſuvian, always exhibits 24 trapezoidal faces, found in volcanic productions; is not fuſible by fire like the red. *Anal.* Silex 55. Argill 89. Lime 6. *Bergman.*

p. *Tourmalin*, of a deep brown, hyacinth, or green colour; does not ſtrike fire with ſteel; can be ſcratched by a knife; cryſtallized in ſix or nine-ſided priſms, longitudinally ſtreaked, and terminating in three-ſided pyramids with pentagonal faces. Is electric when heated. *Anal.* Silex 37. Argill 29. Lime 15. Iron 9. *Bergman.*

q. *Schorl*, of various colours, red, black, greyiſh white, but generally green; ſomewhat tranſparent; ſcratches glaſs, and moſt varieties ſtrike fire with ſteel; cryſtallized in ſix or nine-ſided priſms, with three-ſided pyramids. *Anal.* Silex 52. Argill 37. Lime 5. Magneſia 3. Iron 3. *Chaptal.*

r. *Thumerstein*, Violet ſchorl, vitreous appearance, more or leſs tranſparent; cuts glaſs, and ſtrikes fire with ſteel; found in flat rhomboidal cryſtals, with the two oppoſite edges a little truncated; the ſurface of the ſides ſtreaked, and the ſurface of truncation ſmooth and ſhining. *Anal.* Silex 52. Argill 25. Lime 9. Iron 9. And ſome manganeſe. *Klaproth.*

ſ. *Schorlite*, cryſtallized, indeterminate. *Anal.* Silex 50. Argill 50. *Klaproth.*

t. *Rubellite*, cryſtallized in diverging ſtriated priſms,

prisms, with trihedral summits. *Anal.* Silex 57. Argill 35. Oxyde of iron and manganese 5. *Bindheim.*

u. *Amethyst*, crystallized in hexahedral prisms, with corresponding pyramids; of various degrees of transparency, and colour. *Anal.* Silex 30. Argill 60. Lime 22. Iron 1,66. *Achard.*

v. *Quartz*, crystallized in double hexahedral pyramids, with or without an intermediate prism; of various shapes; strikes fire with steel; is soluble in fluoric, but not in nitric or muriatic acid; *Anal.* Silex 93. Argill 6. Lime 1. *Bergman.*

w. *Prase*, of a leek green colour; crystallized in six-sided prisms, acuminated by six faces, proceeding from the sides of the prisms.

x. *Elastic Quartz*, amorphous, angular. *Anal.* Silex 0,965. Argill 0,025. Iron 0,01. *Klaproth.*

y. *Obsidian*, Iceland Agate, of a greyish or blackish colour; found in irregular shaped pieces. *Anal.* Silex 69. Argill 22. Iron 0,009. *Bergman.*

z. *Calcedony*, Cornelian, Agate, called Calcedon, from the name of a place; usually found in solid masses, of a globular, kidney-like, or stalactical figure, or in the state of pebbles; sometimes filiform, tabular or cellular; of a greyish, yellowish, white, or milk white colour. *Anal.* Silex 84. Argill 16. *Bergman.*

a a. *Chrysoprase*, of a green, or greeniſh white colour; found in ſolid maſſes, and breaks with an even ſurface, having a dull appearance; does not ſtrike fire with ſteel. *Anal.* Silex 0,96. Oxyde of Nickel 0,01. Lime 0,0083. Argill 0,0083. Oxyde of iron 0,0083. *Klaproth.*

b b. *Hyalite*, amorphous, lamellar, compact, colour pure white, fracture even, lamellated, ſometimes conchoidal. *Anal.* 57. Argill 18. Lime 15. And a very little iron. *Link.*

c c. *Opal*, Semi-opal, hard and ſemi-tranſparent; reflects the light in various hues; found of a milk white, bluiſh white and greeniſh colour, rarely purple; takes a fine poliſh, but is not ſo hard as to ſtrike fire with ſteel. *Anal.* Silex 98,75. Argill 0,01. Oxyde of iron 0,01. *Klaproth.* Of this claſs are *Hydrophanes*, which become tranſparent by placing them in water.

d d. *Pitchſtone*, cryſtallized in hexahedral priſms with trihedral pyramids; it's various colours are white, yellow, brown, greeniſh, ſeldom tranſparent; ſcratches glaſs. *Anal.* Silex 73. Argill 18. Iron 00,58. *Wiegleb.*

e e. *Cats-eye*, amorphous, compact, colour greyiſh white, yellowiſh or reddiſh brown, or ſtriped with theſe colours, particularly when poliſhed. Found in blunt or rounded fragments, and comes from Ceylon.

f f. *Flint*, ſtrikes fire with ſteel, generally of a yellowiſh

yellowiſh ſmoke grey, or black grey colour; cryſtallized in double trihedral pyramids; in maſſes of various ſizes, in ſtrata of chalk; when broken, exhibits a conchoidal ſurface, and the fragments are ſharp edged, but of an indeterminate ſhape; ſemi-tranſparent. *Anal.* Silex 60. Argill 18. Lime 2. *Wiegleb.*

g g. *Hornstone, Siliceous Schistus,* or *Basanite, Hornslate. Petrosilex.* Cryſtallized in hexahedral priſms, with or without pyramids, in double trihedral pyramids, in cubes; amorphous, compact; flaty; colour various, yellowiſh white, bluiſh, pearl, green inclining to red, browniſh, roſe colour, blackiſh brown. *Anal. Hornstone.* Silex 72. Argill 22, Carbonate of lime 6. *Kirwan. Basanite.* Silex 73. Lime 10. Magneſia 0,046. Iron 3. Coal 5. *Weigleb. Hornslate.* Silex 73. Argill 24. Iron 3. *Wiegleb.*

h h. *Jasper, Egyptian Pebble,* found in irregular hexahedral priſms; amorphous, in large maſſes and nodules, and of all colours. *Anal.* Silex 54. Argill 30. Iron 16. *Delametherie.*

i i. *Porcelanite,* Porcelain Jaſper, of a pearl grey, or lavender blue colour; found in compact layers. It's ſurface, when newly broken, is a little ſkinny and gibbous.

k k. *Heliotropium, Bloodstone,* of a leek green colour, with blood-red ſpots or veins; is ſemi-tranſparent; exhibits a conchoidal ſurface when broken.

It may be obſerved that Agates do not form a diſtinct ſpecies of ſtone, but conſiſt of quartz, cryſtal, hornſtone, flint, calcedony, amethyſt, jaſper, cornelian, heliotropium, jade, &c. in various combinations, and may be exhibited in the following manner, from *Kirwan*.

The clear pellucid white denotes - -	cryſtal or quartz.
The yellowiſh white -	amethyſt, calcedony;
The greyiſh white -	quartz;
The milk white -	jaſper, amethyſt;
The yellowiſh grey -	quartz, flint;
The ſmoke grey -	calcedony, flint;
The pearl grey -	calcedony;
The greeniſh grey -	calcedony, jaſper;
The bluiſh grey -	calcedony.
The honey yellow -	flint, jaſper;
The yellow - -	jaſper;
The ochre yellow -	calcedony;
The orange yellow -	calcedony;
The yellowiſh brown	cornelian;
The reddiſh brown -	hornſtone, cornelian;
The fleſh red - -	cornelian;
The blood red -	jaſper, cornelian;
The brick red -	jaſper.
The violet blue -	amethyſt;
The browniſh green -	heliotropium;
The leek green, and feeling ſomewhat greaſy -	jade.

11. *Feltspar*, *Adularia*, or Moonſtone, cryſtallized in tetrahedral priſms, truncated obliquely, and

and their varieties; tranſparent; ſhining lamellæ, placed one over the other, often in an irregular manner; but it's figure approaches always either to the cubical or rhomboidal. *Anal.* Silex 62. Argill 17. Lime 6,5. Baroſelenite 2. Magneſia 6. Iron 1,4. *Weſtrumb.*

m m. *Common Feltſpar*, of a pale fleſh colour, yellowiſh, grey, milk white; rarely of a vivid green or bluiſh colour; opaque, compact, ſtrikes fire with ſteel. *Anal.* Silex, argill, with lime and magneſia, or baryte. *Kirwan.* Silex 62,83. Alumine 17,002. Lime 3. Oxyde of iron 1. Loſs. 16,015. *Vauquelin.*

n n. *Labrador Feltſpar*, amorphous, foliated, of various hues, light or deep grey, but when held in certain directions to the light, reflects lazuli-blue, green, and lemon yellow, coppery or violet colour, the ſurface ſhining, texture foliated, fragments rhomboidal. *Anal.* Silex 69. Argill 13. Sulphate of lime 12. Oxyde of copper 0,7. Oxyde of iron 0,04. *Bindheim.*

o o. *Argentine Feltſpar*, cryſtallized, as common feltſpar; amorphous, foliated. *Anal.* Silex 46. Argill 36. Oxyde of iron 16. *Dodun.*

p p. *Staurolite*, cryſtallized in tetrahedral priſms, with tetrahedral pyramids, either ſingle, or croſſing each other at right angles. *Anal.* Silex 44. Argill 20. Baryte 20. Water 16. *Weſtrumb.*

q q. *Lapis*

q q. *Lapis Lazuli*, is opaque, compact, and never found cryſtallized; amorphous, ſeldom pure; of a ſky-blue colour, which it retains in the fire for a long time, but at length becomes brown. *Anal.* Silex, lime, ſulphate of lime, and iron. *Margraff.*

r r. *Prehnite*, ſo called by Werner, after Capt. Prehn, who brought it firſt to Europe from the Cape in 1783. Cryſtallized in tetrahedral priſms; indeterminate; amorphous, foliated; ſemi-tranſparent and brittle; ſurface ſhining, with a lamellated, or fibrous texture; of a greeniſh grey colour. *Anal.* Silex 44. Argill 30. Lime 18. Iron 5. Water and air 2. *Klaproth.*

f f. *Adelite*, of particular ſhapes, tuberous. *Anal.* Silex 62 to 69. Argill 18 to 20. Lime 8 to 16. Water 3 to 4. *Bergman.*

t t. *Zeolite*, opaque, ſeldom ſemi-tranſparent; hard, but rarely ſo hard as to ſtrike fire with ſteel; of a pale green, ſilver white and honey colour, ſometimes of a bluiſh or coppery hue; cryſtallized in cubes; it is found compact, fibrous, radiated, lamellated, uniform, ſtalactical, in drops, and of a capillary ſhape, in which it is very beautiful; in ſix-ſided and flat priſms. *Anal.* Silex 50. Argill 20. Lime 8. Water 22. *Pelletier.*

u u. *Siliceous Spar*, cryſtallized in tetrahedral, and in hexahedral priſms. *Anal.* Silex 61,1. Lime 21,7. Argill 6,6. Magneſia 5. Oxyde of iron 1,3. Water 33. *Bindheim.*

v v. *Rose*

v v. *Rose Spar*, Red Stone, cryſtallized, indeterminate; amorphous, thick foliated.

C. *ADAMANTINE EARTH*, or CORUNDUM, diſcovered by Klaproth, in adamantine or diamond ſpar.

a. *Adamantine Spar*, of a greyiſh colour, inclining to greeniſh white, chocolate brown, &c. cuts glaſs, and ſcratches other gems; ſtrikes fire with ſteel; is not affected by ſulphuric, nitric, or marine acid; exhibits ſix-ſided ſhort rhomboidal priſms, rounded on the top, of a lamellated texture, ſhining in certain directions. *Anal.* Adamantine 68. Silex 31,5. Iron and nickel 00,05, *Klaproth.*

	Corundum of China.		of Peninſula of India.
Argillaceous earth	89,50	—	84,0
Siliceous earth	5,50	—	6,50
Oxyde of iron	1,25	—	7,50
Loſs —	3,75	—	2,0
	100,0		100,0

Chs. Greville.

H. *JARGON EARTH*, diſcovered by Klaproth, in the ſtone called jargon, or zirgon, of Ceylon.

a. *Jargon*, cryſtallized in ſhort tetrahedral priſms, with tetrahedral pyramids; indeterminate, in ſmall grains, and ſmall flat pebbles; ſcratches glaſs;

glaſs: bears a greater heat than the diamond before it is conſumed. *Anal.* Jargon earth 68. Silex 31. Iron and nickel 5. *Klaproth.*

I. *SIDNEIAN EARTH*, diſcovered by Wedgewood; it is contained in a compound ſubſtance from Sidney Cove in South Wales; conſiſting of fine white ſand, ſome colourleſs mica, a few black particles reſembling black lead, and a white argillaceous earth, and from which the new earth is extracted by the marine acid.

a. *Sidneia*, amorphous, looſe. *Wedgewood.*

II. MIXED.

A. CALCAREOUS. Calcareous earth, in it's moſt common ſtate, is called limeſtone, or ſpar; it is in this ſtate that it is combined with a peculiar acid, expellable by heat, and hence called aërial acid, fixed air, or carbonic acid; it effervesces with acids in general: it's more mixed ſtate is now conſidered.

a. *Marl*, Argillo-Calcite, ſemi-indurated or earthy form, of a yellowiſh grey, or yellowiſh white colour; looſe texture, ſtrongly effervesces with acids; in water ſoon falls into powder. In a ſtony or indurated ſtate, fracture earthy, ſometimes ſplintery, or conchoidal, frequently ſlaty; colour various, yellow, grey, brown or bluiſh. *Anal.* Carbonate of lime 60 to 80. Remainder argill; and ſilex. *Kirwan.* In it's ſtony or indurated

durated ſtate, colour yellow, grey, brown, or bluiſh; texture ſplintery or conchoidal, frequently ſlaty; penetrated by bitumen, ſulphur, and pyrites. *Kirwan.*

b. *Limestone with argyllite.*

c. *Siliceous Limestone.*

d. *Ferruginous Limestone.*

e. *Gypsum with Calcareous Spar*; Gypſum is frequently penetrated with this ſpar, ſwineſtone, ſandſtone, and perhaps with ſtrontian; which cauſe a change in it's hardneſs and gravity, and of colour from blackiſh to whitiſh grey.

f. *Gypsum with Marl;* ſometimes diſguiſed with iron; colour reddiſh brown, and reddiſh black; effervesces with acids; gives a red ſtreak.

B. MAGNESIAN.

a. *Calciferous Asbestinite.*

b. *Steatite with Argill.*

c. *Serpentine with Hornblende.*

d. *Siliciferous Potstone;* bluiſh black, mixed with white; it's fracture uneven, partly ſplintery, partly ſlaty; the quartz in many parts viſible in the veins.

c. *Ferruginous Steatite.*

C. ARGILLACEOUS.

a. *Calciferous Argillite;* colour dark, blackiſh, or bluiſh grey, or grey with black or deep blue blotches, or reddiſh or yellowiſh grey; more rarely greeniſh grey; fracture ſlaty, but of ſingle laminæ, ſplintery, or conchoidal.

b. *Talcose Argillite;* colour whitiſh, or bluiſh, or greeniſh grey; often inveſted with foliated ſteatites; found in large maſſes; fracture ſlaty, curved.

c. *Siliciferous Argillite;* penetrated with ſiliceous ſchiſtus, ſand, jaſper, baſanite, or quartz; colour dark, blackiſh, or bluiſh grey, rarely greeniſh grey; fracture ſlaty, but of ſingle laminæ; ſplintery, or conchoidal.

d. *Ferruginous Argillite*; penetrated with calces of iron, of a bluiſh brown colour; externally, it has ſome luſtre; internally, none; ſomewhat of a ſtriated, as well as of a ſlaty appearance.

e. *Hornblende with Garnet;* of a dark greeniſh red, or reddiſh dark green, according to the proportion of garnet; fracture earthy, or fine ſplintery.

f. *Hornblende Slate with Talc or Mica*; colour greeniſh grey; fracture ſlaty, and thin, the laminæ not eaſily ſeparable; found in great plenty at

at Holyhead, and has for the moſt part thick layers of quartz intercepted between it's laminæ.

g. *Hornblende Slate with Quartz;* colour ochre yellow, with black ſhining ſtreaks of hornblende; fracture ſchiſtoſe; feels ſandy; reſembles gneiſs; the yellow colour ſeems to proceed from a ferruginous quartz.

h. *Trap with Hornblende;* colour bluiſh, or greeniſh black; fracture fine ſplintery; contains ſome grains of magnetic ironſtone.

i. *Trap with Mullen;* colour greyiſh black, with a ſhade of red, fracture uneven and fine ſplintery.

k. *Trap with Krag;* colour greyiſh black, with a ſhade of red; fracture uneven and fine ſplintery.

l. *Siliciferous Trap;* colour blackiſh grey, or dark iron grey, with numerous rounded white ſpecks, as ſmall as the point of a pin; it is alſo full of rifts, and theſe exhibit a bluiſh or reddiſh ferruginous illinition; fracture uneven and ſplintery.

m. *Ferruginous Clays;* ſome are ſo penetrated with calces of iron, as ſcarcely to indicate a flaty texture; colour bluiſh brown; externally ſome luſtre; internally none; readily imbibes water.

n. *Mullen with Asbestinite;* colour of a reddiſh grey when firſt broken, with very little luſtre; fracture uneven and earthy.

D. SILICEOUS.

a. *Earthy Quartz;* ſtructure looſe as that of moſs, which it reſembles; exceedingly brittle, and light as pumice; colour white, greyiſh, and dark.

b. *Ferruginous Quartz;* hard, fracture ſplintery; colour brown, or browniſh purple; ſometimes ſchiſtoſe, ſometimes of the texture of moſs.

c. *Earthy Quartz with Actinolite*; greeniſh grey colour, with yellowiſh green lumps, and ſpecks and blotches of red; fracture ſplintery and uneven.

d. *Earthy Hornstone*, reſembles earthy quartz, but of a finer and cloſer grain; colour reddiſh grey mixed with green, browniſh or blackiſh; ſtructure thick ſlaty; fracture ſplintery.

e. *Ferruginous Hornstone;* hard and cloſe grained, colour reddiſh brown; gives a pale red ſtreak.

f. *Siliceous Schistus with Limestone;* of a dark yellowiſh grey colour; fracture ſplintery; the grain very fine and cloſe.

g. *Siliceous Schistus with Argillite;* colour light bluiſh grey or purpliſh grey; fracture fine ſplintery; in the groſs, ſlaty.

h. *Siliceous*

h. *Siliceous Schistus with Mullen;* colour bluiſh grey, it's ſurface often ſtained reddiſh or yellowiſh brown; fracture fine, ſplintery, and uneven.

i. *Pitchstone with Opal;* of a pearl grey, reſembling porcelanite; penetrated with lumps of opal; brittle; imbibes water ſlowly; decrepitates ſtrongly by heat, and then whitens, and remains in a looſe powder.

III. Aggregated.

A. CALCAREOUS.

a. *Calcareous Sandstone*; cryſtallized in rhomboids; amorphous; colour generally white or grey, or yellowiſh white; ſurface rough; fracture earthy or flaty; often ſpangled with mica. *Anal.* Carbonate of lime 37,5. Silex 62,5. *Lassone.*

b. *Calcareous Breccia*; conſiſting of fragments of marble in a calcareous cement.

B. MAGNESIAN.

a. *Potstone Porphyry*; amorphous, undulatingly foliated; greeniſh, reddiſh, or yellowiſh grey; or ſpeckled earth; red or leek green; containing potſtone and feltſpar.

b. *Serpentine Porphyry*; amorphous, compact; dark or olive green, grey, or reddiſh; fracture ſplintery; found near Florence, containing ſerpentine and feltſpar.

C. ARGIL-

C. ARGILLACEOUS.

a. *Argillaceous Sandstone*; amorphous, ſlaty, and compact; does not effervesce with acids; often ſhot through with mica; make mill-ſtones, filtering-ſtones, and coarſe whetſtones; compoſed of argillaceous cement, with fragments of quartz, feltſpar, and flint.

b. *Ruddle Stone*; amorphous, ſlaty, and compact; containing argillaceous cement, with quartz, ſiliceous ſchiſtus, or hornſtone and argillite.

c. *Argillaceous Porphyry*; amorphous; contains feltſpar in indurated clay; hornblende, trap, wakken, mullen, krag, or argillite.

d. *Amigdaloid;* amorphous; containing rounded maſſes of calcedony, agate, zeolite, calcareous ſpar, lithormaga, ſteatite, green earth, &c. in an argillaceous baſis.

e. *Schistose Mica;* amorphous, ſlaty; contains mica and quartz.

D. SILICEOUS.

a. *Granite;* amorphous, compact, ſlaty; chiefly compoſed of quartz, feltſpar, and mica; the compact parts generally irregularly mixed, and of various ſizes.

b. *Sienite;* amorphous, compact, ſlaty; compoſed of quartz, feltſpar, and hornblende; or of quartz,

quartz, feltſpar, hornblende, and mica; ſometimes found ſlaty.

c. *Granatine;* compoſed of quartz, feltſpar, ſchorl; quartz, feltſpar, garnet; &c. &c. &c. *Saussure.*

d. *Granitell;* or Grunſten; amorphous; compoſed of quartz, feltſpar; quartz, ſchorl; quartz, mica, &c. *Kirwan.*

e. *Granalite*; amorphous; compoſed of quartz, feltſpar, mica, ſchorl; quartz, feltſpar, mica, ſteatite, &c.

f. *Gneiss;* amorphous; ſlaty, fibrous, lamellated; contains quartz, feltſpar, and mica. *Werner. Saussure.*

g. *Siliceous Porphyry;* amorphous; compact, ſlaty; conſiſting of cryſtals of feltſpar in a baſis of jaſper, hornſtone, pitchſtone, obſidian, ſiliceous ſchiſtus, ſchiſtoſe hornblende or feltſpar.

h. *Pudding Stone*; amorphous; conſiſting of rounded pebbles in a ſiliceous cement.

i. *Siliceous Sandstone;* amorphous; the grains often ſlightly compacted together, and eaſily breaking into ſand; of this kind is *Salindre*, conſiſting of calcareous grains, inſerted in a ſiliceous cement.

k. *Siliceous Breccia;* amorphous; conſiſting of angular fragments of ſiliceous ſtones in a ſiliceous cement.

CLASS

CLASS III.

METALS.

THE moſt ponderous of all mineral bodies, fuſible, but reſuming their original properties when cold, even after calcination, by the addition of inflammable matter, or oxygen gas: in their pureſt metallic ſtate, they poſſeſs neither taſte nor ſmell. There are 23 metals with which we are acquainted, of theſe 8 are conſidered as ductile, or entire metals; and 15 as fragile or ſemi-metals; of the former are, 1. Platina. 2. Gold. 3. Quickſilver or Mercury. 4. Silver. 5. Lead. 6. Copper. 7. Iron. 8. Tin. Of the fragile are, 1. Biſmuth. 2. Nickel. 3. Arſenic. 4. Cobalt. 5. Zinc. 6. Antimony. 7. Manganeſe. 8. Scheele, Wolfram, or Tungſten. 9. Uranite. 10. Molybdena. 11. Menachanite. 12. Sylvanite. 13. Titanite. 14. Chrome. 15. Tellurium.

From recent experiments made on the Carbonates of Barytes, Magneſia, and Chalk, three new metals are ſuppoſed to exiſt, which Mr. Tondi, the diſcoverer, has named *Barbonum*, *Austrum*, and *Parthenum*; for which Mr. Kerr would ſubſtitute the names of *Barytum*, *Magnesium*, and *Calcum*,

Calcum, as more conſiſtent with the new chemical nomenclature; but further experiments are requiſite to determine the truth of thoſe already made, as they are doubted by ſome eminent chemiſts.

I. Ductile.

A. PLATINA, Plata Silver, or Platina de Pinto; found only in Spaniſh America in ſmall grains among the gold mines there; introduced into England in the year 1749. It is uſually united with iron, and in it's crude ſtate is 18 times heavier than water, and when pure is heavier than gold, it's ſpecific gravity being $21\frac{1}{2}$. With a ſtrong white heat it's parts will adhere together by hammering. The beſt ſolvent is aqua regia, compoſed of equal parts of the nitrous and marine acids.

B. GOLD is a yellow metal, 19 times heavier than water; unalterable by air, or fire; ſoluble in aqua regia, compoſed of three parts of nitrous and one of marine acid, and may be precipitated by various ſubſtances; lime and magneſia precipitate it in the form of a yellow powder; the nut-gall precipitates it of a reddiſh colour; the volatile alkali of a brown, yellow, or orange colour, known by the name of fulminating gold, and this powder weighs a fourth more than the gold made uſe of; liver of ſulphur precipitates

it, by the alkali uniting with the acid, when it falls down combined with the ſulphur; lead, iron, and ſilver precipitate it of a deep and dull purple colour; copper and iron throw it down in it's metallic ſtate. A plate of tin immerſed in a ſolution of gold, affords a purple powder, called the purple powder of Caſſius, which is uſed to paint in enamel. Moſt metals unite with gold in fuſion, hence it is generally found alloyed with ſilver, copper, iron, or other metals.

a. *Native Gold*; found in it's metallic ſtate, cryſtallized with ſilver, copper, or iron.

b. *Grey Ore*; combined with ſulphur, antimony, arſenic, lead, iron, and ſilver. *Born.*

c. *White Ore*; combined with ſilver, biſmuth, and ſulphur.

The largeſt quantities of gold are brought from the Brazils and the Spaniſh Weſt Indies. This metal is found alſo in Hungary, Tranſylvania, and in many other parts of Europe, in red, yellow, black, or iron-coloured ſands; it is met with likewiſe in ſome rivers, as the Tagus, Ganges, Rhine, Saale, Niger, Danube, &c. called river, waſh-gold, or gold-duſt.

C. QUICKSILVER.

QUICKSILVER, or Mercury; the moſt fluid of all metals, not taking a ſolid ſtate until cooled to

to the 29th degree below 0 on Fahrenheit's thermometer, when it becomes malleable; it is 14 times heavier than water, eaſily diviſible, and evaporates in a heat below ignition; rapidly ſoluble in the nitrous acid. Mercury is found in Hungary, Tranſylvania, Carinthia, Bohemia, France, Spain, Sweden, Peru, and probably in the Eaſt Indies and Japan.

a. *Native Quickſilver*; fluid, or interſperſed; ſometimes called virgin mercury.

b. *Amalgamized Mercury*; alloyed with ſilver; found on Moſchellandſberg and Stahlberg in Deuxponts, near Sahlberg in Sweden, near Zlana in Hungary, chiefly on a grey indurated clay.

c. *Cinnabar*; combined with ſulphur, in various forms, ſcaly, granular, indeterminate.

d. *Hepatic Ore*; Mercury combined with ſulphuret of potaſh or ſoda.

e. *Cupreous Mercury*; Quickſilver combined with copper, chiefly on lapis ollaris and quartz. Found in the mines near Moſchellandſberg.

D. SILVER.

SILVER, when pure, is 11 times heavier than water; the whiteſt of all metals; harder, but leſs malleable than gold; ſoluble in the nitrous and vitriolic acids. Silver is found in all countries, but moſt plentifully in Peru and Potoſi.

a. *Native Silver*; found generally of about 16 carats ſtandard.

b. *Arſenical Silver*; cryſtallized with arſenic and iron.

c. *Horn Silver*; reſembling reſin in colour, containing muriatic acid, ſulphuric acid, iron, argill, and lime.

d. *Vitreous Silver*, or Glaſs ſilver ore, of a dark colour like lead ore, contains ſulphur, is ductile.

e. *Brittle Vitreous Ore*, contains iron, antimony, ſulphur, copper and arſenic.

f. *Red Silver Ore*; a brittle red-coloured ore, containing regulus of antimony and ſulphur, and ſulphuric acid. The ſilver is in proportion of 62 to 100. *Klaproth.*

g. *White Silver Ore*; combined with lead, antimony, ſulphur, iron, ſilex, argill; of a light grey colour, and of a dull ſteel-grained texture.

E. LEAD.

LEAD is above 11 times heavier than water, unſonorous, malleable, and very fuſible, ſoluble in all acids and alkaline ſolutions. It yields an oxyde more vitrifiable than the oxyde of any other metal, and a glaſs of a topaz-yellow colour.

a. *Native*

a. *Native Lead*; found in Poland, Silesia, near Karthen, and in Monmouthshire; also in Carinthia, in Vivarais.

(a.) 2. *Native Oxyde of Lead*; crystallized in various forms; compact, or pulverulent. *Anal.* Lead 36. Oxygen 37. Iron 24. Argill 2. *Macquart.*

(a.) 3. *Molybdate of Lead*; crystallized in rectangular tables and their modifications.

b. *Lead Amalgam*; or lead combined with mercury.

c. *Red Lead Ore*, or Ferruginous oxyde of lead; resembling arsenic or realgar; crystallized, shining, striated on the surface; semi-transparent.

d. *White Opaque Lead Ore*, or Native White Lead; of a yellowish or greyish colour. Found at Blegberg.

e. *White Carbonate of Lead*, { Chalk of lead. / Spathose lead. / Mephite of lead.

Spathose lead ore, White transparent lead ore, or White lead spar; colour white or colourless, usually mixed with calcareous earth or clay. Found near Freiburg, Marienberg, in Siberia, Bohemia, Hungary, Tipperary in Ireland, and lead hills in Scotland. *Anal.* Lead 80. Carbonic acid 16, with some lime and argill. *Westrumb.*

f. *Black*

f. *Black Lead Spar*; of a yellowiſh green, or verdegris colour; of a truncated hexahedron form; found on the lead-hills.

g. *Brown Lead Ore*; of a reddiſh brown colour, in four-ſided priſms; found in Tipperary, alſo at Zſchopau.

h. *Native Glass of Lead*; depoſited upon calcareous ſpar, reſembling mica; found at Bergmanſtorff, near Zellerfeld, on the Harz.

i. *Antimonial Lead Ore*; of an iron grey colour. Found at Sahlberg in Sweden, in Siberia; at Lautenthal on the Harz; and in Hungary. *Anal.* Lead 40,50. Antimony 8,16. With ſome ſilver.

k. *Galena*; Sulphuret of Lead, or Sulphurated lead ore; one of the moſt common lead ores, combined with ſulphur, ſometimes ſilver, iron and antimony; it's matrix is generally heavy ſpar, fluor, quartz, coal, ſpathoſe iron ore, ſchiſtus and gneiſs. Found at Bleyſtedt in Bohemia, Freiberg in Saxony; Siberia, Sahlberg in Sweden, and in Derbyſhire. *Anal.* Lead 77. Sulphur 20. Silver 1. *Kirwan.*

l. *Pyritical Lead Ore*; combined with martial pyrites.

m. *Sulphate of Lead*, Vitriol of Lead; found in

in octohedrons, and their modifications; of a yellowiſh brown colour; matrix upon a ferruginous quartz or ſchiſtus. Found in Angleſea, Wales; and near Strontian, Scotland.

n. *Phoſphorated Lead Ore*; lead combined with phoſphoric acid; ſhining, ſemi-tranſparent; found near Freiberg, Zſchoapau, Johanngeorgenſtadt, Zellerfeld, and in the lead hills, Scotland.

o. *Yellow Lead Ore*; or lead combined with acid of Tungſten; yellow or orange coloured lamellated, ſemi tranſparent; is the molybdate of lead of Klaproth, who found it combined with the molybdenic acid. It is found at Bleiberg in Carinthia, near Villach, near Zellerfeld, and in the lead hills in Scotland.

F. COPPER.

COPPER, when pure, is near nine times heavier than water; the moſt ſonorous of all metals, diſſolves in all acids and alkaline ſolutions, oils and water. The leaſt quantity of this metal, in ſolution, turns blue by the addition of volatile alkali: united with calamine, it forms braſs; with tin, bell-metal.

a. *Native Copper*; malleable, fibrous, fuſible, generally found adhering to other foſſil ſubſtances; cryſtallized in cubes, octohedrons, compact, laminar, granular.

b. *Native*

b. *Native Oxyde of Copper;* found in compact lumps, ſprinkled in the ſtate of ſmall particles, of a hyacinth colour. Sometimes reſembling brown pitch, or of a ſteel grey, and browniſh red. Cryſtallized in cubes and octohedrons.

c. *Pitch Copper Ore*, hyacinth red; little ſhining, ſoft, compact, pulverulent; containing oxyde of copper with oxyde of iron. *Born.*

d. *Carbonate of Copper*; Malachite, ſky blue, moſtly pulverulent, cryſtallized in rectangular octohedrons and various; ſtalactitical. Found in Tyrol. *Anal.* Copper 73. Carbonic acid 26. *Fontana.*

e. *Arseniate of Copper*; of a deep olive, or emerald colour, ſemi-tranſparent, cryſtallized in various modifications. *Anal.* Copper 73. Carbonic acid 26. *Fontana.*

	Synonymous Names.
f. *Sulphate of Copper*,	Vitriol of Cyprus. Blue Vitriol. Vitriol of copper, or of Venus. Blue copperas.

cryſtallized in tetrahedral priſms, and their modifications, ſhining, colour braſs yellow: contains copper, iron, and ſulphur.

g. *Muriate* of Copper*; amorphous, in a ſandy

* Salt formed by the union of the muriatic acid with different baſes.

form.

form. *Anal.* Copper 52. Acid 10. Oxygen 11. Water 12. Sand 11. *Rochefoucauld.*

h. *Sulphuret of Copper*, Vitreous Copper Ore, or Pyrites of Copper; of a lead grey colour, cryſtallized in hexahedral truncated priſms; contains copper with ſulphur; ſometimes a little ſilver or arſenic. Found at Freiberg.

i. *Variegated Copper Ore;* exhibiting various colours; intermixed with copper pyrites and vitreous copper. Found at Freiberg.

k. *Yellow Copper Ore;* cryſtallized in equilateral tetrahedrons, and various; compoſed of copper, iron and ſulphur.

l. *Grey Copper Ore;* cryſtallized in tetrahedrons and their modifications, and in ſix-ſided priſms. *Anal.* Copper 16. Lead 34. Antimony 16. Iron 13. Sulphur 10. Silver 2. Silex 2. *Klaproth.*

m. *White Copper Ore;* ſilver white, brittle; on rubbing emits an arſenical ſmell; compounded of copper, iron, and arſenic. Found near Freiberg.

6. IRON.

IRON is about eight times heavier than water, is attracted by one of it's ores, called the loadſtone; ſoluble in all acids, alkaline ſolutions, water, and air; it's ſolution is turned of a black or

dark purple colour, by galls, and other vegetable aſtringents.

a. *Native Iron;* various, irregular, malleable; of an iron grey colour; found at Kamſdorf, in Siberia; and South America.

b. *Grey Iron Ore*, Magnetic Iron Stone, Emery; ſhining, copper red colour, a little attracted by the magnet; cryſtallized in octohedrons, and in cubes; lamellated, foliated, granular, and various; contains iron united to a ſmall proportion of oxygen.

c. *Hematite;* of a blood red, or ſteel grey colour; found ſolid, reniform, fibrous, ſcaly and various; found at Schneeberg, &c.; contains iron with carbonic acid and argill. *Kirwan.*

d. *Argillaceous Iron Ore*, Baſaltic Iron Ore; reniform, globular, lenticular, granulated, various; found at Bohnerz in Germany; contains iron united to oxygen, carbonic acid and argill, and often phoſphate of iron.

e. *Spathose Iron Ore;* of a greeniſh grey cream colour, or chocolate brown; cryſtallized in rhombs, and in double, three, and ſix-ſided pyramids, foliated; found near Freiberg, Kamſdorf, &c. in Germany. *Anal.* Iron 38. Lime 38. Carbonic acid and manganeſe 24. *Bergman.*

f. *Sulphate*

Synonymous Names.

f. *Sulphate of Iron.* { Martial vitriol. Green vitriol. Vitriol of iron. Green copperas.

a combination of the ſulphuric acid with iron.

g. *Sulphuret of Iron,* Martial Pyrites; cryſtallized in tetrahedrons, and their modifications, in cubes, octohedrons, ſolid, interſperſed, capillary, ſtriated; conſiſts of iron and ſulphur.

Mr. David Muſchet, of the Clyde iron works, divides iron-ſtones into,

1. Argillaceous Iron-ſtone, having clay for it's chief component earth, and this clay comparatively pure and free from ſand.

2. Calcareous Iron-ſtone, poſſeſſing lime for it's chief mixture, and this lime alſo comparatively deſtitute of ſand.

3. Siliceous Iron-ſtone, uniting clay and lime, containing large proportions of ſilex.

4. Iron-ſtone containing nearly the ſame proportions of clay, lime and ſilex.

Beſides theſe are deſcribed, "Primary ores of iron," ſo named in contradiſtinction to ores which appear, like iron-ſtones, to have been formed by a ſecondary agency.

H. TIN.

TIN is a ſilver-coloured gliſtering metal, ſeven times heavier than water; does not vitrify like lead; is malleable and unſonorous; ſoluble in aqua regia, vitriolic, muriatic, nitric and acetous acids; yields a white oxyde that impairs the tranſparency of glaſs, and converts it into enamel. Found in Cornwall and Bohemia.

a. *Native Tin:* pure native tin is ſo very rare, that it's real exiſtence has been doubted.

(a.) 2. *Native Oxyde*, Spathoſe tin ore; tin combined with oxygen and iron; is found in various modifications.

b. *Brown Tin-ſtone and Spar;* conſiſts of calx of tin, calx of iron, and acid of tungſten.

c. *Wood Tin*, Stream Tin, or Corniſh Tin Ore; found only in Cornwall, in ſmall globular or reniform pieces, ſometimes of a fibrous or radiated texture; containing tin, with oxygen and iron.

d. *Tin Pyrites*, Sulphuriſed Tin, or Sulphuret of Tin; contains tin, ſulphur, copper and iron, beſides it's matrix; it is diſtinguiſhed by it's ſulphureous ſmell when heated.

II. Fragile.

A. BISMUTH.

BISMUTH is of a white yellowiſh colour, and laminated texture; above nine times heavier than water, very fuſible, ſoluble in the ſulphuric muriatic and nitric acids, and particularly in the laſt, and when diſſolved in it, is precipitable by a mere dilution with pure water; the precipitate is white. Found plentifully in Saxony, Bohemia, Sweden.

a. *Native Bismuth,* of a ſilver white colour, inclining to reddiſh. It's matrix is red jaſper, petroſilex, quartz, heavy ſpar, and cobalt ores. Found cryſtallized in equilateral triangular laminæ; indeterminate; in ſmall particles, foliated. Separated from it's matrix by ſimple fuſion; being itſelf eaſily fuſible.

b. *Native Calx, or Oxyde of Bismuth;* without luſtre; friable, looſe, or compact; of an earthy appearance; emitting no ſulphureous ſmell when ignited; ſoluble in nitric acid; of a yellowiſh grey, greeniſh, and ſtraw yellow grey.

c. *Sulphurised Bismuth,* or Sulphuret of Biſmuth; of a lead grey and bluiſh grey colour; ſometimes variegated; amorphous, ſtriated, foliated, looſe and diſperſed. *Anal.* Biſmuth 60. Sulphur 40. *Sage.*

d. *Martial*

d. *Martial sulphurated Bismuth.* Of a radiated texture; yellowiſh grey colour. Found near Gillabet in Norway.

e. *Arsenicated Bismuthic Ore;* of a yellowiſh white colour; brilliant luſtre; emits a garlick ſmell when ignited; conſiſts of biſmuth, arſenic, and ſulphur. Found at Schneeberg in ferruginous jaſper, accompanied with cobalt ore.

B. NICKEL.

Found generally in a metallic ſtate, more rarely in the ſtate of calx. When free from heterogeneous ſubſtances, is of a grey reddiſh white, or fleſh colour; when broken, has a ſtrong luſtre; of a fine-grained, compact texture; has a little ductility; is ſoluble in ſulphuric, muriatic, and particularly nitric acid.

a. *Native Nickel;* copper-red colour; exhibits a conchoidal ſurface when broken; cryſtallized in rhomboidal tables; amorphous, foliated. It's matrix is calcareous ſpar, and heavy ſpar. *Anal.* Nickel alloyed by iron. *Born.*

b. *Native Calx or Oxyde of Nickel;* colour of a pale green, or bluiſh green; has an earthy appearance; is friable; ſometimes ſhining. *Klaproth* found it mixed with ſiliceous, argillaceous, magneſian, and calcareous earth.

c. *Kupfer Nickel,* copper-red colour; deeper than

than the purified metal; of particular ſhapes; amorphous, granular, compact. *Anal.* Nickel united to iron, arſenic, cobalt, and ſulphur. *Bergman.*

C. ARSENIC.

ARSENIC is about eight times heavier than water, appears in plates of a bluiſh grey colour, may be diſcovered by evaporating it upon red-hot coal or iron, by means of which a garlic ſmell is emitted. The fumes depoſit a white coating on a plate of copper. It is not acted upon by water, but readily by nitric acid. Phlogiſticated alkali dropped into it's ſolution, produces a green precipitate.

a. *Native Arsenic*, Teſtaceous Arſenic; arſenic alloyed by iron. It is generally found in cobalt mines, in Saxony, Bohemia, Norway.

b. *Native Oxyde of Arsenic*, White Oxyde of Arſenic; this is ſcarce, of a whitiſh colour, but expoſed to heat becomes blackiſh. It is ſoluble in water aſſiſted by heat; alſo in nitric acid, leſs ſo in muriatic and vitriolic acid. Found in Hungary, on the Hartz, in Bohemia, and Tranſylvania.

c. *Sulphuret of Arsenic*, Sulphuriſed Arſenic, Realgar, yellow Orpiment; of a yellow brimſtone colour, inclining to orange; it's texture uſually lamellar, and ſo ſoft as to be cut with a knife, and a little

a little flexible; often of a beautiful brilliant luſtre, which is improved by the ſcarlet red ſhades. When reduced to powder, it is uſed as a pigment. Found in Hungary, and other places. *Anal.* Arſenic 84,90. Sulphur 16,10. *Kirwan.*

d. *Red Sulphurized Arſenic*, or Ruby Arſenic; of an aurora colour; contains a larger proportion of ſulphur than the preceding; common in China, where it is made into vaſes, pagodas and other ornamental works. Red arſenic is commonly found near volcanos, as at the Solfatara near Naples, &c.; it's matrix is quartz, heavy ſpar, and ferruginous clay.

e. *Miſpickel*, Pyritical Arſenical Ore, Arſenical Mundick; arſenic and iron mineraliſed by ſulphur; ſometimes combined with ſilver; it is found in cubes, rhomboidal, four-ſided, truncated priſms, ſometimes terminating in dihedral ſummits, with triangular plans; octohedral; the ſurface is generally ſtriated. When encloſed in a cloſe veſſel, it ſublimes and forms orpiment, leaving the iron behind. It's matrix is ſpathoſe iron ore, fluor, quartz, blende. Found in Saxony, Bohemia, Tuſcany, and on the Hartz.

D. COBALT.

COBALT is ſeven times heavier than water, of a roſe-white colour, brittle and eaſily reducible to powder; is a little moveable by the magnet, probably from ſome iron it contains; difficult of fuſion; when fuſed with ſand and potaſh, it forms ſmalts,

ſmalts, uſed for blueing clothes. Is ſoluble in nitric acid without heat, and in vitriolic acid aſſiſted by heat; likewiſe in the marine and nitro-muriatic acid, which diluted with water, is uſed for ſympathetic ink.

a. *Grey Cobalt Ore*, Arſenicated Cobalt; chiefly compoſed of cobalt and arſenic; of a fine-grained, compact texture; of a ſteel-grey colour. Found in Saxony, Bohemia, Norway, in a matrix of red heavy ſpar; calcareous quartz.

b. *White Cobalt Ore*, conſiſts of cobalt and iron, mineraliſed by ſulphur and arſenic, of a tin-white colour, of a granular and lamellated texture. Found in Norway and Sweden.

c. *Sulphuret of Cobalt*; contains no arſenic or iron; when heated it emits a ſulphureous ſmell; found in cubical cryſtals in Upper Hungary, generally upon quartz.

d. *Native Oxyde of Cobalt*; Black Oxyde or Calx of Cobalt; this is the pureſt kind of the calciform cobalt ores; when heated, it emits no ſulphureous or arſenical vapours. Found in Thuringia, Saxony, Tyrol. Of the oxydes of cobalt there are various ſpecies, as *Brown earthy Oxyde, Yellow Oxyde, Green Oxyde, Red Oxyde, of Cobalt.*

N. ZINC.

ZINC is about 7 times heavier than water; it is ſoluble in the nitric, ſulphuric, and muriatic acids, and produces by the two laſt hydrogen gas or inflammable air; it burns with a bluiſh green flame when expoſed to red heat, and ſublimes in the ſtate of a white light ſubſtance; it precipitates lead and other metals, in a metallic ſtate, from their ſolution. When in an oxyde ſtate, is recovered by charcoal in a cloſe veſſel. It is found in various parts of Europe, never in a perfect metallic ſtate, but generally in the ſtate of calx or oxydated; frequently alſo combined with iron and ſulphur. It is of a fibrous texture, and of a bluiſh grey colour.

a. *Native Oxyde of Zinc*, calamine, calx of zinc. *Anal.* Oxyde of zinc, 84. Oxyde of iron 3. Silex 12. Argill 1. *Bergman.*

Synonymous Names.

b. *Carbonate of Zinc*, { Chalk of Zinc. / Aërated Zinc. / Mephite of Zinc.

Oxyde or Calx of Zinc, combined with, or mineraliſed by fixed air or carbonic acid. Is perfectly ſoluble in ſulphuric acid, without emitting heat; the fixed air is diſcovered by the effervescence which takes place when diſſolved in the ſulphuric acid, and the zinc combining with the acid, forms

c. *Sulphate*

Synonymous Names.

c. *Sulphate of Zinc*, { Vitriol of Zinc. / White Vitriol. / Vitriol of Goſlar. / White Copperas.

ſoluble in water; formed from blende, acidified by the abſorption of oxygen, and thus producing the ſulphuric acid, which afterwards combining with the calx of zinc, compoſes the Sulphate or Vitriol of Zinc. *Anal.* Zinc 20. Sulphuric acid 40. Water 40. *Bergman.*

d. *Blende*, or Sulphurized Zinc; combined with ſulphur, iron, copper, ſilex. It is of various colours, ſometimes reſembling galena, and hence called pſeudo-galena; when of a blackiſh colour, it is called *blackjack*. *Blende*, in the German, ſignifies blinding or deceitful. *Anal.* Zinc 52. Sulphur 26. Iron 8. Copper 4. Silex 6. Water 4. *Bergman.*

O. ANTIMONY.

ANTIMONY is about 7 times heavier than water; ſoluble in aqua regia, and precipitated by pure water; ſcarcely ſoluble in ſulphuric or nitric acid; with muriatic acid it forms butter of antimony.

It is found in nature in the metallic ſtate; in the ſtate of calx, mineraliſed by arſenic, but moſt generally with ſulphur, called crude antimony. It is of a bluiſh colour, of a lamellated and radiated texture.

 a. *Native*

a. *Native Antimony*, uſually mixed with iron and arſenic. It contains ſo large a proportion of the latter, that by fuſion with ſulphur, the product reſembles realgar, or red arſenic; found in Dauphiny, Sweden, Saxony.

b. *Arsenicated Antimony*, Native arſenical Antimony; of a white brilliant luſtre, and ſcaly texture; when heated, it emits only arſenical vapours. *Anal.* Antimony alloyed by arſenic.

c. *Muriate of Antimony*, White antimonial Ore; combined with muriatic acid, of a greyiſh white colour, found in oblong, rectangular, four-ſided laminæ. It is ſcarce; found near Braunſdorf, Freiberg, Bohemia; generally accompanied by red blende, galena, grey antimony.

d. *Red Antimonial Ore*; conſiſting of antimony, ſulphur, and arſenic; of a light crimſon red colour, in hexahedral, priſmatic, capillary cryſtals; found in the antimonial mines of Bohemia, Hungary, and Tranſylvania.

e. *Sulphuret of Antimony*; Sulphuriſed Antimony; of a beautiful appearance of the rainbow colours; filaments long, in hexahedral priſms, with obtuſe tetrahedral priſms; it is extremely ſcarce; it was found in a certain antimonial mine of Hungary, in quartz and heavy ſpar. *Anal.* Antimony 74. Sulphur 26. *Bergman.*

f. *Plumose Antimonial Ore*, or Argentiferous antimonial

antimonial Ore; antimony combined with iron, arſenic, ſulphur, and ſometimes ſilver; of a lead-grey colour; cryſtallized in ſlender priſmatic needles. It is found near Freiberg, in Hungary and Tuſcany.

P. MANGANESE.

MANGANESE is about 7 times heavier than water; ſoluble in marine acid; the ſolution appears brown; by addition of marine acid to manganeſe, is produced an elaſtic fluid or gas, called ſuroxygenated muriatic gas, or dephlogiſticated muriatic acid air. When expoſed to red heat in a cloſe veſſel, pure air is produced. It is uſually found in the ſtate of calx, or combined with oxygen. Next to iron and gold, it is the moſt frequently diffuſed metallic ſubſtance through the earth, and even in vegetables. Soluble in acids, and precipitated in a white powder by alkali.

a. *Native Manganese*; found in the ſtate of greyiſh-white ſmall globules, which, on expoſure to the atmoſphere, fall to a black powder.

b. *Native Oxyde of Manganese*, White oxydated Manganeſe; cryſtallized in tetrahedral rhomboidal priſms; found in Hungary, Tranſylvania, mixed with ſilex, and conſtitutes the matrix of the auriferous ores. *Anal.* Oxyde of Manganeſe 43. Oxyde of Iron 43. Lead $4\frac{1}{2}$. Mica 5. *Wedgewood.*

c. *Black*

c. *Black calciform Manganese*, Black-Wad; of a greyiſh, or iron-black colour; found in Könitz, Piedmont, Carinthia, Thuringia, England, and Ireland.

d. *Siliceous Ore of Manganese*; of a ſteel-grey colour, and lamellated texture; combined with oxyde of manganeſe 35. Silex 55. Iron 5; and argill 5.

Q. SCHEELE.

SCHEELE, *Wolfram*, or *Tungsten*; of a greyiſh white, or yellowiſh colour; granulated, friable; is never found in the metallic ſtate, but alone in the ſtate of calx, combined chiefly with calcareous earth, manganeſe and iron; it's oxyde is of an acid nature, of a yellow colour, and enters into combination with all metals; it is ſoluble in the ſulphuric, nitric, muriatic, and nitro-muriatic acids, and is converted by them into an oxyde, which combines alſo with alkalis, but is precipitated again from them by nitric acid. With the muriatic acid, the oxyde aſſumes a blue colour.

a. *Tunſtate of Lime**, Sparry Tunſtate of Lime, Tungſten; combined with Tungſtenic acid 44; and lime 56; of a lamellar texture; greyiſh white colour; cryſtallized in angular octohedrons, or double tetrahedral pyramids; is 6 times hea-

* Salt formed by the combination of the Tunſtic acid with different baſes.

vier

vier than water; digeſted with muriatic acid, it turns yellow; found in Bohemia and the tin mines in Cornwall.

b. *Wolfram*, Manganeſeous Wolfram; combined with tungſtenic acid 64; oxyde of manganeſe 22; oxyde of iron 13; ſilex and tin 2; of a dark black colour; lamellated, brittle, cryſtallized in compreſſed hexahedral priſms, terminated with tetrahedral pyramids; found in England, Siberia, Bohemia, Saxony.

R. URANITE.

URANITE, or Uranium; is of a deep-grey colour, with a ſlight luſtre; above 6 times heavier than water; is ſoft enough to be cut with a knife; ſoluble in nitric, ſulphuric, and marine acids; is not precipitated by zinc; phlogiſticated alkali added to the ſolution, produces a deep red precipitate. Firſt ſeparated in 1790, by Klaproth, from the mineral called Pechblende. Found in a ſtate of calx or oxyde, and mineraliſed by ſulphur.

a. *Carbonate of Uranite*; Calciform or Oxyde of Uranite, or Calcolite. The earthy oxyde of uranite is a variety of an earthy texture; of a brimſtone yellow; it is found ſolid, diſperſed through, and depoſited upon other ores. The variety called ſpathoſe uranite, is of a graſs-green colour, and generally found cryſtallized in cubes, and in hexahedral priſms; conſiſts of uranium, carbonic

carbonic acid, and a little copper; found in Bohemia.

b. *Sulphuret of Uranite*; Sulphurated Uranite, Pechblende; of an iron colour, brittle texture; emits a ſulphureous ſmell expoſed to heat.

S. MOLYBDENA.

MOLYBDENA, is of a ſteel-grey colour, very brittle; 6 times heavier than water; is only found in nature combined with ſulphur. It melts eaſily; is volatile by moderate heat, effervesces with alkali; emits a ſulphureous ſmell when treated with the blowpipe.

a. *Sulphurised Molybdena*, or Molybdena mineraliſed by ſulphur; conſiſting of the molybdic acid and ſulphur; it's matrix is feltſpar, lithomarge, and quartz; found in Iceland, Sweden, Spain, Saxony, France, Siberia; frequently in rocks containing wolfram and tin ores.

T. MENACHANITE.

MENACHANITE, diſcovered in the valley of Menachan, in Cornwall, by Mr. Gregor; and ſuppoſed to conſtitute a new metallic ſubſtance.

a. *Native Menachanite*; Menachanite alloyed by iron; found amorphous, or in grains, and in irregular forms; of a grey, or dark colour; ponderous, and attractable by the magnet.

U. SYL-

U. SYLVANITE.

This ſemi-metal, diſcovered by *Facebay*, is different from all other known metallic ſubſtances; and is called Sylvanite from it's being found in Tranſylvania; of a metallic luſtre; fracture broad, or granularly imbedded; of a dark-grey or white colour; reſembles regulus of antimony, or ſulphurated biſmuth; evaporates by continued heat; amalgamates with mercury by ſimple trituration; and detonates with nitre; ſoluble in aqua regia, and in concentrated vitriolic acid in cold, or a low digeſting heat, and the ſolution is crimſon red; but by the affuſion of water, or by a ſtrong heat it is precipitated. Muller found it to contain a ſmall proportion of arſenic and of nickel, and alſo of gold. Bergman found it to contain a little zinc; but theſe mixtures ſeem to be merely caſual. *Kirwan.*

V. TITANITE.

A ſemi-metal, diſcovered by Klaproth, found at Rhonitz, and at Bainick in Hungary; of a brownish-red colour; fracture foliated, unequal and conchoidal; cryſtallized in right-angled quadrangular priſms, longitudinally ſtreaked and furrowed, often acicular, and ſeated on ſchiſtoſe mica alternating with quartz. Expoſed to a moderate heat, it remains unaltered, but in a cool crucible it burſts into angular fragments, loſes it's luſtre and colour, and becomes pale brown; inſoluble in aqua regia, vitriolic, nitrous, or marine acid.

a. *Calcareo Siliceous Ore*; found near Paſſau; diſcovered and deſcribed by Profeſſor Hunger; of a reddiſh, yellowiſh, or blackiſh-brown colour; found maſſive or diſſeminated; but more frequently cryſtallized in obtuſe-angled tetrahedral cryſtals; found ſeated on, or inhering in gneiſs or granite; foliated or ſtriated; cuts glaſs. *Kirwan. Anal.* Silex 35. Calcareous earth 33. Titanitic calx 33. *Klaproth.*

W. CHROME.

CHROME, ſo called from it's property of colouring the combinations into which it enters; diſcovered by Klaproth in the red lead of Siberia. Vauquelin has made many experiments upon this ſubſtance, but it's place in mineralogy is not yet fully eſtabliſhed.

X. TELLURIUM.

Klaproth has chemically analyzed the auriferous mine, known as the mine of white gold (Weiſe goldertz) aurum paradoxum, metallum vel aurum problematicum; found in the mine called Mariahulf, in the mountain of Fatzbay near Zalethna, in Tranſylvania; and in this mineral has found a metal different from every other yet known.

It is of a whitiſh greyiſh colour; friable and lamellar; in cooling after being heated, it takes a cryſtallized ſurface; very hard of fuſion. Placed on a heated body, it burns with a bright flame, of a blue colour; it amalgamates with mercury.

It's

It's ſolution in the nitrous acid is clear and colourleſs; when it is concentrated, it exhibits pointed needle-like cryſtals.

The pure alkali precipitates from the acid ſolutions of it, an oxyde of a white colour, ſoluble in all the acids.

The ore of white gold of Fatzbay, aurum vel metallum problematicum, contains: *Anal.* Metal of Tellurium 925,5. Iron 72,0. Gold 2,5. Total 1000.

The graphic gold of Offenbanya, contains Tellurium 60. Gold 30. Silver 10. Total 100.

The mineral known under the name of the *Yellow Ore of Nagyag*, contains Tellurium 45. Gold 27. Lead 19,5. Silver 8,5. Sulphur ſcarcely an atom. Total 100,0.

The *Foliated grey Gold of Nagyag*, contains Lead 50. Tellurium 33. Silver 8,5. Sulphur 7,5. Silver and Copper 1. Total 100,0.

CLASS IV.

INFLAMMABLES.

INFLAMMATION, or combuſtion, conſiſts in the fixation and abſorption of vital air by combuſtible bodies, and in the decompoſition of atmoſpheric air by theſe bodies. As vital air is a gas, and many combuſtible bodies, by abſorbing, fix it, and make it take a ſolid form; hence vital air, when thus precipitated, will loſe the caloric to which it owed it's elaſtic fluidity; and hence comes that free caloric or heat, which is evolved during combuſtion.

I. Aeriform.

A. *Hydrogen*,

Is one of the component principles of water; and in combination with caloric and light, forms hydrogen gas.

a. *Pure Hydrogen*, exiſts always in a ſtate of gas. It is a conſtituent of water; where it forms one ſixth. It is diſtinguiſhed from all other gaſes, on

on account of it's being inflammable; hence it is called inflammable air. See §. V. page

b. *Sulphurated Hydrogen, Hepatic Gas,* is obtained from livers of ſulphur, or ſulphures, by decompoſing them with acids. Sulphur combines with hydrogen gas, called ſulphuret, or hydrogen gas, or ſulphurated hydrogen gas, or hepatic air.

II. Simple bituminous Substances.

Bituminous ſubſtances are not of mineral origin, but have been formed from certain principles of ſubſtances belonging to the organiſed kingdoms of nature; which, after the loſs of animal and vegetable life, have ſuffered conſiderable changes, by long contact and union with mineral bodies; and may hence be regarded as forming a part of the mineral ſyſtem.

a. *Naptha,* is a light, thin, often colourleſs oil, highly odoriferous and inflammable, which in ſome parts of Perſia and Italy is found upon the ſurface of ſprings and lakes, and iſſuing from argillaceous ſtones; does not combine with water or ſpirit of wine; when burned, leaves a black ſoot; acids condenſe it and render it reſinous: when firſt expoſed to the air, it becomes yellow; afterwards, and in like proportion, it thickens, and paſſes into

b. *Petroleum, or Rock Oil;* and ſeems chiefly to differ from Naptha in being mixed with heterogeneous

geneous matter; it is found in various parts of Europe trickling from the fiſſures of rocks. It has a greaſy feel, is tranſparent, or ſemi-tranſparent, of a reddiſh or blackiſh-brown colour; by air it becomes like tar, and then is called

c. *Mountain or Mineral Tar, Barbadoes Tar,* Bitumen Petroleum tardefluens. It is viſcid; of a reddiſh or blackiſh-brown or black; when burned it emits a diſagreeable bituminous ſmell, and by expoſure to the air, it paſſes into

d. *Mountain or Mineral Pitch, Bitumen Maltha;* which is more impure than petroleum; it has the conſiſtence of honey; is of a browniſh-black colour; frequently mixed with much earthy matter, reſembles common pitch; when heated emits a ſtrong unpleaſant odour, like the former ſubſtance; in cold weather it may be broken, and exhibits internally a gloſſy luſtre; when warmed it ſoftens and poſſeſſes ſome tenacity. It is however ſuſceptible of a ſuperior degree of induration, and then becomes *asphaltum.*

e. *Asphaltum,* or *Bitumen of Judea,* has derived it's name from the Dead Sea of Judea, upon the ſurface of which it is found, and upon the ſhore. It is met with alſo in other countries. It is ſmooth, of a black colour, and ſhining conchoidal fracture.

III. COM-

III. Compound bituminous Substances.

Mr. Hatchett, whoſe diviſion of bituminous ſubſtances I have chiefly followed, is of opinion that the progreſſive changes of naptha into petroleum, mineral tar, mineral pitch, and aſphaltum, is cauſed by the gradual diſſipation of part of the hydrogen of the bitumen, and the conſequent diſengagement of carbon; and although the diviſion of the ſimple bituminous ſubſtances terminates in aſphaltum, nature appears to have glided on by an uninterrupted chain, which connects the ſimple bitumens with the compounds.

a. *Jet*, is a compact black body; harder and leſs brittle than aſphaltum; may be formed into trinkets; it breaks with a conchoidal fracture, and the internal luſtre is gloſſy. It has no odour, unleſs when heated, and it then reſembles aſphaltum.

b. *Coal*, of which carbon is a conſtituent principle, is found in various parts of Europe in ſtrata beneath the ſurface of the earth; it is foliated, brittle, black, and glittering. Beſides carbon, it's conſtituent parts are, petroleum, maltha, and aſphaltum, combined more or leſs with other earthy particles, and, frequently with ſulphur, and the remains of vegetable matter.

c. *Mineral Elaſtic Gum;* diſcovered in 1786, near

near Caſtleton, in Derbyſhire; reſembles, in colour and elaſticity, cahout-chou, or Indian rubber; formed from naphtha or petroleum; there are many varieties; ſome in the ſtate of aſphaltum, by melting loſe their elaſtic property, and a quantity of gas is diſengaged; and the ſubſtance remaining reſembles mineral pitch, or mineral tar.

B. AMBER is one of the pureſt bitumens, conſiſting almoſt entirely of a volatile oil, which receives it's conſiſtence from a peculiar acid (ſuccinic) and of a few earthy and coaly particles.

Amber has been chiefly procured on the ſhores of the Baltic. The yellow amber of Kœnigſberg has been lately diſcovered by digging. It is raiſed 200 feet from the ſea, where many ſhafts are ſunk, one of which is near 100 feet in depth. The amber does not run in veins, but is found in nodules, in a matrix of charcoal, below which there are ſtrata of ſand. In the charcoal there are ſometimes little threads of amber. This ſeems to ſhew that the origin was volcanic, and that the amber was accumulated by ſublimation.

a. *Honeyſtone;* cryſtallized in aluminiform octohedrals; found in Thuringia, between beds of bituminous wood, and is ſometimes near an inch in diameter; it is cryſtallized in great maſſes, in a double pyramidal form, with four ſides; it bears a great reſemblance to amber, but does not, like that ſubſtance, become electric by rubbing; by Mr. Abich's analyſis it appears, that 50 parts of this foſſil are compoſed of 8 parts of the carbonate of

of alum, 2 of carbon, $1\frac{1}{2}$ of oxyde of iron, 20 of carbonic acid, 14 of the water of cryſtallization, emitting the ſmell of bitter almonds, and $2\frac{1}{4}$ of ether; with a ſmall portion of the benzoic acid. In conſequence of the incombuſtibility of the honey-ſtone, he propoſes to exclude it from the claſs of combuſtible bodies, and to place it among that of aluminous earths.

b. *Common Amber* is found in ſeveral countries of Europe beneath the ſurface of the earth, among clay, ſand, and the iron bog-ore, but moſt abundantly in the ſea, and particularly on the ſhore of the Baltic; found in maſſes, irregular, tranſparent or opaque, of a white, yellow, or brown colour; red and green amber being ſcarce. Taſteleſs; of a faint odour when rubbed, and becomes electric; inſoluble in water.

C. MINERAL TALLOW, or MUMIA; found in 1736, in the ſea, on the coaſt of Finland; alſo in ſome rocky parts of Perſia, mixed with petroleum. It is perfectly white, of the conſiſtence of tallow, but more brittle though as greaſy; it burns with a blue flame, and a ſmell of greaſe, leaving a black viſcid matter, which is more difficultly conſumed.

D. SULPHUR is a ſimple body, which, during combuſtion, or, in other words, during it's combination with oxygen, produces ſulphuric acid. Sulphur is found near the craters of volcanos, and the ſources of ſome mineral waters.

a. *Common, or Native Sulphur*, is a yellowiſh ſubſtance, odorate, electric, tranſparent, and octohedral; opaque and priſmatic; fuſible, and ſubject to different combuſtions; one is ſlow, with a bluiſh flame, and yields ſulphureous acid; the other rapid, with a white flame, and yields ſulphuric acid.

b. *Volcanic Sulphur;* amorphous, granular, compact.

E. PLUMBAGO, or *Carbure of Iron*; or iron combined with a large portion of carbon.

a. *Common Plumbago; Graphite;* this ſubſtance has not yet been produced by art, but nature contains it in conſiderable abundance; ſuffers no change by water or atmoſpheric air, nor expoſed to fire in cloſe veſſels; but burns ſlowly, if heated, in contact with air; amorphous, fibrous and ſlaty; of a deep iron black colour; little ſhining, metallic luſtre, ſtains the fingers, ſoapy to the touch.

b. *Anthracolith;* conſiſting of Carbon 90. Argill 5. Iron 3. *Born.*

VOLCANIC

VOLCANIC PRODUCTIONS.

I. CINDERS.

A. Loose.

a. *Ashes*, of a browniſh or reddiſh grey colour; conſiſtence looſe and duſty; very light and ſubtle, and ſmooth to the touch; ſlowly diffuſible in water, and when wet, ſomewhat ductile; effervesce ſlightly with acids; and contain about half their weight of argill, a ſmall proportion of earth, calx, magneſia, and iron, and the remainder is ſiliceous.

b. *Sand;* conſiſting of minute hard grains, that readily ſink in water; generally fragments of lava and ſcoriæ, together with olivia, garnets, feltſpar, and ſchorl.

B. Coherent.

a. *Puzzolana;* colour reddiſh brown, grey, or greyiſh black; that of Naples grey; that of Civita Vecchia, reddiſh brown; ſurface rough, uneven, and of a baked appearance; pieces of various ſizes, from the ſize of a nut to that of an

egg; fracture uneven, or earthy, and porous; commonly filled with particles of pumice, quartz, ſcoriæ, &c.; very brittle; does not effervesce with acids; magnetic before it is heated, but not after; contains ſilex, argill, and iron.

b. *Trass*, or *Terras*; reſembles puzzolana; though often the product of pſeudo volcanos, or external fires; colour greyiſh brown, or yellowiſh; ſurface rough and porous; fracture earthy, rarely lamellar; found principally near Andernach, in the vicinity of the Rhine; alſo near Frankfort, Cologne, Pleith, &c.; and there called Duffſtein; containing fragments of argillite and baſaltine, often branches of trees half cleared, and impreſſions of leaves, with mica, iron ore, &c.

c. *Tafa*; a kind of puzzolana formed by nature into a mortar; colour brown, or reddiſh brown, brick red, or ſpeckled with various colours; fracture earthy; contains ſand, ſcoriæ, fragments of lava, limeſtone and pumice; often baſaltines and Veſuvian; commonly magnetic, and not eaſily decompoſed by the action of the ore.

d. *Pumice*; colour grey, rarely brown, or blackened by fuliginous fumes; ſurface rough and fibrous, with elongated pores; ſometimes fibres not diſcernible; fracture ſtriated and open, uneven and ſplintery; fragments oblong, obtuſe, and irregular; brittle in a high degree; fuſible at 130. o. into a grey flagg.

e. *Piperino*;

c. *Piperino*; a concretion of volcanic aſhes, and is ſaid to be the ſubſtance that covers Pompeia; colour grey or reddiſh brown; fracture earthy; contains fragments of white marble, feltſpar, mica, garnets, ſcoriæ, gypſum, ſchorl, granite, &c. ſometimes magnetic; if preſerved from moiſture, it hardens in the air.

II. LAVA.

Any matter that has iſſued out of a volcano in a liquified ſtate, or that has accompanied or been enveloped in ſuch liquified matter. *Kirwan.*

a. *Cellular;* colour brown, or greyiſh black; ſurface unequal, rough, and full of cavities; affects the magnetic needle.

b. *Compact;* earthy ſubſtance, or matter, which, after having been fuſed, but not vitrified, becomes, on cooling, compact, cloſe, and ſolid.

III. VITREOUS LAVAS.

a. *Glass*; ſuch matter as has been expoſed to the greateſt heat, and from the compoſition, moſt ſuſceptible of vitrification in a moderate heat; colour black, green, yellow, or greyiſh white; found in detached maſſes; either ſpongy or compact fracture.

b. *Enamel*; reſembling vitreous lavas, is the imperfect vitrification called enamel; colour white,

white, grey, or brown, often with round ſpots, of a different colour from the ground; ſometimes contains cryſtals of ſchorl, or feltſpar, not completely melted; fracture rather more granular than that of glaſs.

c. *Scoriæ*; compounded of iron and ſtony matter, and owe their fuſion to ſulphurated iron; they are tumefied and expanded by the ſulphureous vapours; colour generally black; ſometimes reddiſh from the calcination of the iron; ſurface rough, and uneven; texture cavernous; the cavities larger and more irregular than thoſe of cellular lava, but never fibrous.

d. *Slaggs*, reſemble the droſs of forges, generally red and heavy.

THE

NATURALIST's and TRAVELLER's COMPANION, &c.

PART the SECOND.

ALTHOUGH it may he admitted with peculiar honour to the preſent age, that the knowledge of natural hiſtory and of ſcience in general has been of late conſiderably enlarged; yet as the objects of human inquiry are numberleſs, and frequently diſperſed in diſtant parts of the globe, as well as complicated in their hiſtory, the ſentiments of an ancient philoſopher may be adopted even at this day with propriety: "Multa etenim "ſunt quæ eſſe audivimus, qualia autem ſint ig-"noremus! Quamque multa venientis ævi popu-"lus, ignota nobis, ſciet*!"

At the ſame time if we reflect upon the foregoing ſuggeſtion, reſpecting the amazing progreſs made in natural hiſtory within the ſpace of a few years, we may find ſufficient inducement to perſevere in purſuits ſo worthy of a rational mind.

* Seneca.

It

It would render natural hiſtory much more pleaſing, as well as greatly tend to it's progreſs, were the limits of our knowledge therein preciſely aſcertained, that travellers and curious perſons, who have little leiſure for reading, might not only be informed of what is already diſcovered, but alſo of what is ſtill doubted, or unknown; by which means their inquiries would be better directed, and more conducive to real information and uſeful diſcovery.

SECT.

SECT I.

Observations and Queries respecting Learning, Antiquities, religious Rites, polite Arts, &c.

Ingenuous arts, where they an entrance find,
Soften the manners, and ſubdue the mind*.

1. THE alphabets of the various nations, their pronunciation and numeric value, with their numeric figures, if different from the letters of the alphabet, and books written in each language, eſpecially grammars, dictionaries, &c. with the dates of each when written, merit the inveſtigation of the curious; likewiſe the materials uſed for writing, and their preparation, as the methods of making ink, paper, and pens, and of ſizing and gluing the paper; the art of printing, and the contrivances for doing it.

2. Manuſcripts, in good preſervation, of the Hebrew Bible, or parts thereof, particularly if upwards of 300 years old.

3. Books containing the religious principles of any nation or people, and which uſually are written in a dialect different from that which ſuch people now ſpeak, or in a poetical, high metaphorical ſtyle, and therefore underſtood by few only, and for the moſt part kept very ſecret;

* ——Ingenuas didiciſſe fideliter artes
Emollit mores, nec ſinit eſſe feros.

OVID. Pont.

amongſt theſe we may enumerate the Chartah Bhade Shattah of Bramah, the Chartah Bhade and Aughtorrah Bhade Shaſtah, the Vedam*, the ſacred books of the Perſees, written in the ancient Perſian dialects, called the Zend, and Pehlvi†; the Koran of Mahomet; the ſacred books of the Mendæans or Sabaites, at or near Baſſora in the Perſian gulph; the voluminous ſacred books of the Lamas in Nexpal and Thibet, called Khangiur, and it's myſtic part, termed Riuté; the ſacred books of the prieſts in Pegu and Siam, and others of ſimilar tendency. Theſe would be ſtill more valuable, could Engliſh, Latin, or French tranſlations be added.

4. Deſcriptions of the manners, cuſtoms, feaſts, and religious ceremonies of the reſpective nations; the architecture both exterior and interior of their temples, religious, public, and private buildings; the figures, names, genealogies and ranks of their divinities and idols; their ſacred and domeſtic utenſils, the caſts and ranks of people, the learning and religious tenets of each nation, to all which explanatory drawings would be required. What nations uſe circumciſion, and what are the advantages derived from ſuch a cuſtom, or diſadvantages from the omiſſion of it? Is circumciſion ever extended to the females, and in what manner is it employed?

* In the peninſula on this ſide the Ganges, the ſacred books of the Bramins are contained in the Vedam, copies of which, in the original Sanſcrit character, would be very valuable.

† The Pehlvi is a more modern dialect.

5. The

5. The tranſlations of the Bible in different languages; and the ſacred books of Chriſtians of various denominations; as the Georgian, Armenian, Perſian, Æthiopian, Coptic, Arabic, and Syriac, eſpecially among the Chriſtians on the Malabar coaſt, and in the iſle of Socatora*.

6. The hiſtory and ſucceſſion of princes, the origin and migration of nations; the government and political conſtitution of each country; the cauſes of the increaſe or decay of power.

7. A relation of the private and domeſtic life of the people; the cuſtoms obſerved at the birth of children; the marriages, ſepulchral rites, and any other circumſtances characteriſing each nation.

8. An account of the aſtronomy and chronology of different nations, whether they obſerve the ſyſtem of the ſeven days of the week, the names for theſe days, with their ſignification. The number, names, and ſignifications of their months; the number of theſe in a year; whether they are uſed to conciliate the moon's and ſun's motion by any intercalation, or a certain cyclus of years; the names for particular ſtars and conſtellations in the zodiac, with their ſignifications; the diſtinctions of the other ſtars from the planets, with the length of their revolutions.

The ſtate in which the art of drawing, carv-

* The Neſtorian Chriſtians formerly had a ſettlement among the Indians on the Malabar coaſt, and were there very much reſpected. Are ſome of theſe ſtill exiſting? Have they any ancient Syriac books?

ing, and engraving in ſtone has been, or now is. Specimens, drawings, or collections of old inſcriptions, engravings, ſeals, gems, ſtatues, carvings, baſſo and alto relievos, and the places where each of the above monuments are found, the ſize, ſubſtance whereon it is worked, &c. the ancient and current coin, with the exact valuation.

SECT.

SECT. II.

Commerce, Manufactures, Arts, Trade, &c.

Jove has the realms of earth in vain
Divided by th' inhabitable main:
If ships profane, with fearless pride,
Bound o'er the inviolable tide*.

FRANCIS.

DESCRIPTIONS and drawings of the looms, tools, machines, &c. employed in manufactures, particularly if simple, ingenious, and gaining time or strongly increasing power, might prove highly beneficial.

2. An account of the planting, gardening, and agriculture of each country; the manure used, the time and labour employed in each branch of business; the price of labour, the implements of agriculture; the kinds and quantities of corn sown in an acre; the quantities reaped in different soils; the proportion of vineyards or pasture lands to arable, and the number of people in a square mile of pasture, arable lands, vineyards, or any other kind of plantation.

3. The kinds of pigments, stains, and dying materials known and used, particularly in China;

* Nequicquam Deus abscidit
Prudens oceano dissociabile
Terras, si tamen impiæ
Non tangenda rates transiliunt vada.
HOR. l. 1. Od. 3.

are

are they mineral or vegetable? the manner of preparing and applying them, with the advantages and disadvantages of each sort compared with ours; particularly the materials, machines and methods employed by the Indians for dying, staining, and printing their chintzes, calicoes, &c.

4. The wood and timber used for ship building; the form and construction of the ships; the wood employed for masts; the succedanea for oakum, ropes, cables, sails, pullies, &c. with the comparative advantages and disadvantages.

5. The means devised for catching quadrupeds, birds, fishes, shells, &c. either for food, or to prevent the increase of such as are noxious to the people or their plantations; are any animals made tame and employed to catch others, or are any methods used for killing or inebriating them?

6. The materials of clothing; if animal skins, the manner of dressing them; if the hair of animals, or the threads of certain insects, the method of spinning, twisting and weaving such substances; if vegetables, how are they cultivated, dressed, spun, and manufactured? The cut and make of the dresses in general, with the advantages and disadvantages of each particular part.

7. The various objects of commerce in general, the growth and manufacture of each article, with the names by which it is known, and it's uses, when designed only for inland trade; the price of labour, and the number of people employed in each department.

8. It is a common opinion, that large quantities of remnants and rags of all kinds of scarlet cloth,

cloth, are yearly carried from England to China, and that the Chineſe extract from them their fine red pigments. If this be true, what methods are employed to extract the colour?

9. It has been obſerved that analogous ſubſtances are moſt proper for dying homogenous bodies*; thus animal ſubſtances are beſt for dying wool and ſilk, becauſe wool and ſilk are animal ſubſtances. A blue dye, made of woad (Iſatis of Linnæus) is found to be full of inſects. Is it not the ſame caſe with indigo? Are not all the laſting dyes made from animal ſubſtances, or of ſuch as contain numerous inſects?

10. The manner in which the beſt indigo is manufactured in the interior parts of Indoſtan, and the plant from which it is made. Is it from the Indigofera or the Anil? Are there any rules to aſcertain when the plant has ſoaked ſufficiently, and how long it ought to be beaten?

11. Is there any linen made of flax or hemp, or what other ſubſtances are ſpun and wove in India beſides cotton? What uſe is made of the yellow or brown cotton taken from the Bombax? Is it manufactured for apparel, and appropriated for a certain order of men, as prieſts or Bramins?

12. Deſcriptions and drawings of the inſtruments and machines employed by the Chineſe and Indians to clean the cotton from the ſeeds.

13. Is only European zaffer from cobalt uſed by the Chineſe for painting their porcelain blue, or have they ſome of their own? If they have,

* Vid. Hiſtoire de l'Academie, an. 1768. art. 11.

what

what name is it known by, and how is it manufactured? It is probably finer than ours, from the richneſs of the old China figures.

14. The preparation of the pickle or catchup called Soya; is it made from the Dolichos Soya Linn.? Is ſalt, wheat or barley added, and in what proportions?

SECT.

SECT. III.

Metereological Observations, Food, Way of Living, Animal Œconomy in general, &c.

> For every man to native cuſtom prone,
> Conforms and models life to that alone.
>
> GOLDSMITH.

IT is always ſatisfactory to have regular meteorological accounts by the aſſiſtance of a good barometer and thermometer; and to obſerve at the ſame time the quarter the wind blows from, and it's degree or violence; the quantity of rain and ſnow by inches; the ſize of hailſtones; the appearance of aërial phenomena, as aurora borealis, or northern lights, fiery globes, halos or bright circles round the ſun and moon; with the effects likewiſe of thunder ſtorms, lightning, &c.

2. The traveller ſhould alſo remark the ſucceſſion of ſeaſons, and the various fruits and productions of each country; the times of ſowing or planting, as well as of harveſt, or of reaping the grain, &c. the budding or flowering of trees, or ſhrubs. The food of the inhabitants, and the preparation of it previous to it's uſe.

3. Some account might be collected of the general prevailing diſeaſes in different ſeaſons, and the cauſes producing the ſame, or the remedies employed for curing them, and the methods in which remedies are adminiſtered.

4. Are any diſeaſes cauſed by the effluvia of certain trees or plants, or is the touching, handling or cutting of trees or plants ever ſuddenly prejudicial to health? Does the effluvia from ſpice trees or the frankincenſe tree prove deleterious?

5. What are the effects or ſymptoms which ariſe from the bite of poiſonous ſnakes, or any reptile or inſect? Has the bite of a ſnake ever been found to have a ſalutary effect in curing a certain previous diſeaſe, or does the bite of one ſnake ever deſtroy the effects of another? Are there any counterpoiſons or antidotes uſual againſt the bite of ſnakes? Is muſk or any ſpecies of the ariſtolochias a remedy which the ſnakes avoid and fly from, or do any of theſe prove lethal to ſnakes?

6. Is the pedra de cobra uſed as an antidote againſt the effects of the bite of any ſnake, and with what ſucceſs? How is this remedy procured and applied?

7. The manner of managing domeſtic animals, whether in health or otherwiſe. The animals which the natives in any country caſtrate, and the effects produced by it; as well as the period when ſuch a cuſtom was firſt introduced.

8. It would deſerve remarking where and in what manner polygamy is introduced, whether perpetual or temporary, and the effects of it upon the manners, religious or civil cuſtoms, population, &c. of ſuch countries. Does polygamy prevent ſome men from procuring wives, or are the women brought from neighbouring nations? Is it

any

any where cuſtomary for one woman to be married to ſeveral huſbands during the life of the firſt?

9. Accurate calculations of births or burials in provinces or towns, and the proportions of males to females, would prove very valuable.

10. Have any buildings or ſhips, furniſhed with electrical rods after Dr. Franklin's method, ever received any injury from lightning?

11. Is the venereal diſeaſe cured without mercury? and if ſo, by what remedies?

12. What diſeaſes attack the workmen employed in different kinds of manufactories?

SECT. IV.

Zoology.

Non ad unam natura formam opus ſuum præſtat; ſed in ipſa varietate ſe jactat.

SENECA Quæſt.

1. IT would greatly tend to improve our knowledge in this department of natural hiſtory, were the following remarks reſpecting quadrupeds to be carefully made; viz. the general times of coupling and of geſtation; how many young are brought forth at a time, and how often during one ſeaſon; at what period of life they become prolific or barren: where their principal reſort and dens are; whether the males aſſiſt the dams in providing food for the young of the carnivorous tribe; how long theſe are under the protection of the old ones; and what age each ſpecies attains?

2. It might be inquired whether any perſon hath ever ſeen elephants in copulation, which has been hitherto denied; it is ſaid that if the wild elephants perceive any body, they immediately begin to rave, and ceaſe not till the curioſity of that perſon has been rewarded with death; and though the Indian princes have kept great numbers of tame elephants of both ſexes, they never could procure a breed from them. What differences are there betwixt the African and Indian

Indian elephants? Is the ſtructure of the grinding teeth equal or flat in all, or have ſome elevated or pointed crowns, ſimilar to thoſe of carnivorous animals? Do elephants ever ſhed their tuſks or teeth, or are they permanent?

3. How many ſpecies of tigers, ſo called, or more properly of leopards, panthers, ounces, &c. are there in India, and what are the ſtated and perpetual marks for diſtinguiſhing each, in different periods of life? What other animals of the feline kind are found in India, with the ſpecific characters of each?

4. Does the ſhakal, or jackal, (canis aureus Lin.) bear any reſemblance to an animal commonly called the croſs-fox (Pennant's Syn. Quadr.) What animal do the Arabian writers call banat-el-auvi?

5. Are armadilloes (daſypodes Lin.) in Aſia or Africa, in what parts, and what are their characteriſtics?

6. When the various ſpecies of the feathered tribe begin to couple or pair ſhould be noticed; when, where, and of what materials each bird builds it's neſt, with the colour, ſize and number of eggs; how long the eggs are in incubation; what the young are fed with, and at what periods they are fledged; with the diſtinctions between male and female birds in different ſeaſons and ages.

7. The migration of birds ſhould not be diſregarded; but their merely diſappearing in one part of the country is not properly a migration, for we frequently find that birds ſhift their place of abode,

at

at certain ſeaſons, on account of ſome palatable food, which may be more plentiful in one part than in another; the croſſing wide ſeas or extenſive continents is underſtood. If any bird be found out at ſea, the ſpecies of bird, the direction of it's flight, the diſtance from land, and the latitude and longitude ſhould be noted.

> ———In figure wedge their way,
> Intelligent of ſeaſons; and ſet forth
> Their airy caravan, high over ſeas
> Flying, and over lands; with mutual wing
> Eaſing their flight: ſo ſteers the prudent crane
> Her annual voyage, borne on winds: the air
> Floats as they paſs, fann'd with unnumber'd plumes*,

In northern climates it would be uſeful to obſerve when ſwallows are firſt ſeen, and when they diſappear; and likewiſe in what climates they have been found, whether in a torpid or active ſtate, with the ſpecies and peculiar characters.

8. What birds or animals are allowed a privilege or immunity from being injured or killed? and what may be the reaſons for the ſame, or the advantages derived therefrom?

9. Some birds of prey are employed in the eaſt by the grandees in hawking and hunting. What means are uſed for teaching ſuch birds, and what are their differences, ſize, figure, plumage, names, ſpecific characters, &c.?

10. What are the character, plumage, food, and œconomy of the Indian ravens (buceros Lin.)

11. Are there any humming birds (trochilus

* Milton.

Lin.)

Lin.) in the Indies or China with a filiform long tongue, conſiſting of two ſemi-cylinders?

12. What are the peculiar diſtinctions of the true wild peacock? Are there any white peacocks in India, and of a ſeparate ſpecies? Is the change in plumage obvious in wild peacocks, or is this the reſult of domeſtication? Do white peacocks breed with grey and green ones, and what is the colour of the young breed?

13. Is there to be found in the Indian ſeas a jaculator fiſh, ſciæna jaculatrix*, different from the chætodon roſtratus, Linn.? or is the faculty of ſhooting at inſects with a drop of water peculiar to theſe fiſh, or common to any others?

14. Has the rajah torpedo, Linn. or the cramp or numb-fiſh, the ſame electric qualities as the gymnotus electricus?

15. As there is yet wanted a good figure of the ſea cow (trichechus manatus Lin.) it would be deſirable to procure a good drawing of it while alive, to have it diſſected, and to obſerve wherein it coincides with, or differs from other animals nearly related to it, as the ſeal, dolphin, &c.

16. The ſeaſons ſhould be noticed, when different ſpecies of fiſh ſpawn, and the rivers, bays, ſhoals, or ſands they reſort to for that purpoſe; what age they attain before they ſpawn, the food they eat, the age they live to as accurately as poſſible, the ſize they acquire, and the latitude wherein they are generally found; the method of catch-

* Philoſophical Tranſactions, Vol. 54. and Vol. 56. t. 8. p. 186.

ing

ing them, and to what uſes alſo they are applied when caught, and whether they are eſteemed wholeſome food or the contrary?

17. Which ſpecies of moth is it, the caterpillar of which in China affords that ſtrong grey kind of ſilk, and how is it manufactured or wove? How are theſe ſilkworms or caterpillars preſerved, fed, and managed? The introduction of ſuch a new ſilk into England would be a uſeful acquiſition, and redeem entomology from the cenſure it is now branded with, of being a mere curioſity void of any real utility.

18. Are there any rattle-ſnakes in the Indies or China, or in any part of the world beſides America?

19. The various bundles of ſwimming ſea weeds ought not to be neglected; for beſides the different kinds of fuci they conſiſt of, they generally contain ſmall crabs, ſhrimps, or other ſubmarine inſects or worms, ſuch as the oniſci, monoculi, ſometimes ſhells and eſcharæ, ſertulariæ, and other corallines.

20. The various animals inhabiting ſhells deſerve inveſtigation, as there are probably new genera yet to be diſcovered: it is not fully aſcertained whether the inhabitant of the paper-nautilus be an animal really belonging to that ſhell, or only uſes it in the ſame manner as ſome crabs do other ſhells; neither is it known whether or no the paper-nautilus animal, or the ſepia, increaſes or enlarges it's ſhell. Indeed the generation, and general œconomy of all the molluſca claſs are very imperfectly known.

21. Experi-

21. Experiments might be tried to aſcertain whether pearls are not ſuch concretions in the ſhells wherein they are found, as the crabſtones (lapides ſeu oculi cancrorum) in the ſhells of the river cray-fiſh, which collect alkaline or calcareous materials from the food of the animal for the formation of the new ſhell, as do the mya and mytilus margaritifer a new layer for the increaſe of their ſhells. Do the ſhells containing the pearls gradually increaſe through the year, or at a certain ſeaſon only?

22. It ſhould be carefully remarked which of the ſhells, eſpecially of the murex and turbo kinds, afford a red or purple juice fit for ſtaining or dying like the purple of the ancients; whether the dye be permanent, and the method of preparing and applying it by ſome Indian nations. D'Ulloa, in his South-American Voyage, Vol. I. page 182, ſpeaks of one of theſe ſhells found near Panama on the Darien Iſthmus; and Janus Plancus, or Giovanni Bianchi, in his work on ſhells, mentions the turbo ſcalaris Linn. or wendeltrap, as affording the violet purple of the Romans.

23. Deſcriptions and drawings of the animals or polypes inhabiting the various corals, corallines, ſponges, echini, or ſea-eggs, aſteriæ or ſtar fiſh, ſepiæ or cuttle fiſh, holothuriæ or ſea-nettles, and all the various molluſca, and the polypes of the tubipores, madrepores, millepores, cellepores, &c. deſerve the attention of a natural hiſtorian, as but few of them are well deſcribed or known.

SECT. V.

Botany.

Itaque iſta quoque naturæ rerum contemplatio, quamvis non faciat medicum, aptiorem tamen medicinæ reddit.

CELSUS.

1. THE ſeeds of almoſt all the Indian plants are worth collecting, they may be preſerved in the manner deſcribed in ſect. III.

2. What trees bear the myrobalans, a drug formerly much in requeſt, but at preſent in little eſteem as a medicine, but which might probably afford ſome uſe in dying? What ſpecies of trees bear the myrobalans bellirica, chebula, citrina, and indica or nigra; whether a kind of phyllanthus as the myrobalanus emblica? The uſe alſo of the ſame in India?

3. Is the aloe-wood or eagle-wood, the calambac, and the agallochus, the ſame or different? The place from whence procured, with the generic and ſpecific characters?

4. Is the orchel or rochel (lichen rocella, Lin.) which is found in Madeira, and uſed to dye red, a kind of lichen or a zoophyte? Is the ſteeping it in putrid urine ſufficient to prepare it?

5. Where does the lignum rhodium grow? Is it a kind of nyctanthes or Indian jaſmine?

6. Of what genus and ſpecies is the tea-wood, of which tea-cheſts are made?

7. What

7. What is the wood tek on the Malabar coaſt, of which the Indian ſhips are built? Is it a fact that it is never attacked by ſea worms (teredo, Lin.)?

8. The various kinds of pulſe, as peaſe, beans, phaſeoli, &c. eſpecially ſuch as are reared at the Cape of Good Hope, and exported to the Indies; the other fruits likewiſe which are cultivated at the Cape, and taken in as refreſhments on board the European ſhips.

9. The different kinds of palm trees, their nature, ſoil, characters, names given by the natives, and the uſes they put them to; or their fruits, leaves, bark, pulp, &c.

10. To what genus and ſpecies does the graſs called Tatack belong? and where does it grow, beſides Madagaſcar, Java, the Malay Iſlands, and the French Iſle de Bourbon? The graſſes in general which thrive in particular countries and climates, with the ſoil and culture, and the kinds of cattle moſt addicted to each.

11. What plant bears the famous Indian nut, which is uſed as a reſtorative, and is immenſely dear, being ſold according to ſome at three thouſand pounds each; the place where it is cultivated, the ſoil it requires, and it's real or imaginary virtues?

12. The Columbo root, called by the Portugueſe Raiz de Mozambique, is a native of the continent of Aſia, but it has been tranſplanted to Columbo in Ceylon, and the Dutch now ſupply all Aſia with it. Is it different from the root of Lopez or

Lopezia; and if ſo, what are the characters of each?

13. A deſcription of the ſmall grains and phaſeoli, with which the Indians on the Coromandel coaſt ſow their fields after the rice harveſt, with the minutiæ of their cultivation, eſpecially the machines employed for watering the grounds.

14. A tree or plant in Cochin China, called Tſai, on being fermented like Indigo, plentifully furniſhes a green-coloured flour, which in dying gives a laſting tincture of a fine emerald; it would be therefore worth inquiring after the method of extracting the colour, and the additional ſubſtances employed in fixing it, and what ſtuffs are beſt fitted to receive the ſame.

15. Are the ſtamina of the pterocarpus draco, Lin. which is called Draco arbor ſiliquoſa populi folio, by Commelin, and Lingoum by Rumph. Amb. 2. t. 70. connected or ſeparate?

16. Which genus of plants does the true ebony belong to? Is it an aſpalathus?

17. Many varieties of rice are found in India, as the red, with red huſks; the little rice, ſmall grained, oblong and tranſparent; the great long rice, with round grains; the dry rice, which grows beſt on a dry ſoil, and requires no watering; and the common rice; are theſe various kinds of rice different ſpecies or varieties only? The culture, characters, and ſpecimens of each, if different, would be neceſſary.

18. There is an elaſtic gum, called Borrachio in Portugueſe, and Kaboutchuck in the language of the natives near Cayenne in South America, of which

which it is ſaid the Chineſe make rings for laſcivious purpoſes, but here uſed by ſurgeons for injecting liquids, and by painters for rubbing out black-lead-pencil marks. Is this gum manufactured in India or China, and from what plant, and in what manner, with the different uſes it may be applied to? Is the plant an Euphorbia or Apocynum?

19. What plants produce gum myrrh and gum ammoniac, and how are they collected?

20. What plant affords the gummi rubrum aſtringens from Gambia?

SECT.

SECT. VI.

Mineralogy.

And join both profit and delight in one*.

CREECH.

1. THE manner of working mines, and the methods employed in getting, breaking, and extracting the ores; the tools and machines employed for each of these purposes, are subjects worthy of inquiry. Is gunpowder ever used to blast the stones or ores? The manner in which the ore or metal is found under ground, whether in perpendicular veins, or in vicinity to them; in horizontal flat strata, loose pieces, or in solid continued bodies; in what kind of stone, and at what depth; the means of carrying off the water when present in the mines. The vapours found in them, whether mephitic and noxious, or inflammable when fire or light comes in contact with them.

2. The manner in which white copper, resembling silver, is manufactured, and the various processes whereby it is done.

3. The operations used in extracting the metals from the respective ores, with specimens and names of the ores, and the places where they are

* Simul et jucunda et idonea dicere vitæ.

HOR.

procured;

procured; the products yielded from the ores by fusion; the fluxes added to promote fusion, or the substances to prevent volatilization, and whatever is subservient towards refining of metals or regulises; with the structures and materials of the ovens, the fuel and quantities of it employed, the time for each operation, and the preparatory cautions, including the picking, pounding, washing, sifting, and ustulating of the ores; and drawings of the various machines and tools used for each purpose.

4. The places from whence the various gems or precious stones are procured, with their prices on the spot; the ground and strata wherein they are found, and the figure or form of each kind before being cut, whether determinate and general, or accidental.

5. The manner of manufacturing those immense quantities of saltpetre annually exported from the East Indies; the soil employed for the lixivium, and the manner of preparing the soil. Are any animal or inflammable substances added to it? By what means is lixivium precipitated? Is an alkali used for that purpose, and how is the alkali procured? Is any use made of the remainder of the lixivium after the precipitation of the saltpetre? Is the lixivium boiled, and inspissated by fire for the crystallization, or by the heat of the sun?

6. If borax be artificial, in what manner and from what substances is it made? If native, in what strata and soil does it lie?

7. How far has the knowledge, value, and use of metals extended amongst nations?

SECT.

SECT. VII.

Directions for taking off Impressions or Casts from Medals and Coins.

——Et cætera penè gemelli.
HORAT. Ep. X.

CHIEFLY owing to the coſt required for purchaſing a cabinet of medals, it has happened that the ſtudy of them has hitherto been confined, comparatively, to a few individuals. Another principal impediment to the cultivation of an acquaintance with them, has ariſen from the difficulty of underſtanding the inſcriptions, for want of a ſufficient knowledge of languages; on which account, in particular, this ſtudy has been condemned by the illiterate as barren and uſeleſs; but ſuch as are acquainted with the advantages which have already reſulted from thoſe *nummi memoriales*, cannot heſitate a moment to aſſiſt in promoting a more general purſuit of the ſubject*.

While Coloſſian ſtatues, and the hardeſt marbles, with their deepeſt inſcriptions, are deſtroyed by accidents, or by time, and paintings finiſhed with the higheſt colours quickly fade, a medal ſhall ſurvive innumerable accidents, and diſcloſe hiſtorical facts a thouſand years after ſtatues are

* Though the ſtudy of medals does not properly belong to natural hiſtory, this ſhort account of taking impreſſions from them, may prove acceptable to ſome travellers.

crumbled

crumbled away; and when nothing but the names of an Apelles or a Praxiteles remain. Does not a ſingle medal, of which we are in poſſeſſion, give us greater light into hiſtory, than the once famous libraries of Alexandria and Pergamos, which are now no more? From theſe, and many other conſiderations, I would willingly contribute my endeavours to render this ſtudy more general, and conſequently more uſeful. I have tried a variety of methods to enable a young medalliſt to collect a cabinet, which may initiate him into the knowledge of medals and coins at a trifling expence.

The method of taking off plaſter of Paris and ſulphur impreſſions, is known to every body: the firſt is too ſoft to preſerve them from injury, and the brittleneſs of ſulphur is a greater objection.

By forming a coat or layer of thin metal over the plaſter of Paris, it would be a conſiderable defence. Tin is the cheapeſt and moſt convenient metal for the purpoſe, as it is ſufficiently flexible, and at the ſame time very much reſembles ſilver. The tin-foil ſhould be of the ſame kind with that uſed for ſilvering looking-glaſſes. It ſhould be laid over the medal or coin intended to be taken off, and then rubbed either with a bruſh, the point of a ſkewer, or a pin, till it has received perfectly the impreſſion of the medal; the tin-foil ſhould now be pared off round the edge of the medal, till it is brought to the ſame circumference: the medal muſt then be reverſed, and the tin-foil will drop off into a chip-box or mold ready to receive it, the concave ſide of the foil, or that which is laid on the face of the medal,

being uppermoſt; upon this pour plaſter of Paris made in the uſual manner, and when dry, the figure may be taken out of the box or mold, with the tin-foil ſticking on the plaſter, the convex-ſide being now uppermoſt again, in which poſition it is to be kept in the cabinet, after it becomes dry. To have an impreſſion very perfect, the thinneſt tin-foil ſhould be made uſe of*.

The impreſſions taken in the foregoing manner almoſt equal ſilver medals in beauty, and are very durable: if the box or mold † be rather larger than the impreſſion of tin-foil, the plaſter, when poured on, runs round it's edges, and forms a kind of white frame, or circular border, round the foil, whence the new-made medal appears the more neat and beautiful. If this tin-foil be gilt with gold-leaf, by means of thin iſinglaſs glue, or boiled linſeed oil, the medal will reſemble gold.

* This method does not in the leaſt injure any medal or coin.

† Chip boxes, uſed by apothecaries, anſwer this purpoſe, and may be eaſily procured. A ſlip of paper wrapped round any circular body with a flat ſurface, is equally convenient.

INDEX.

INDEX.

	Page
ACID, - -	96
..... of Borax, -	96
..... Sulphur,	97
..... in water to discover,	69
Actynolite, -	111
Adamantine earth,	103, 107
..... spar,	127
Adularia, -	124
Aedelite -	126
Aërated acid	45, 85
..... calcareous earth,	104
..... iron,	43, 71
..... lime, -	67
..... magnesia,	60
..... mineral fixed alkali,	100
..... ponderous earth,	107
..... soda,	46, 100
..... vegetable fixed alkali,	98
..... zinc, -	154
Aërial acid, -	96
Agallochus, -	194
Agate, -	124
Agriculture of different countries, -	181
Aigue marine,	118
Air, contents of to discover,	81
Air, contents of,	81
Alkahest of Van Helmont,	98
Alkali in water to discover,	69
Alkaline air, -	83
Alkaline salts, -	46
Aloe wood, -	194
Alphabets of different nations,	177
Alum, -	69, 112,
.... in water, -	43
Alumine, -	103, 112
Aluminous gas, -	83
..... sulphate	69
Amber, - -	168
..... common -	169
Amethyst, -	121
Amianthus, -	111
Ammoniac, -	101
Ammoniacal chalk	101
..... gas,	83
..... nitre,	102
..... salt, secret of Glauber -	102
Amorphous, -	105
Animal earth, -	106
..... œconomy, queries concerning, -	xi
Animals, how domesticated	156
Antimony -	36, 155
..... white ore of,	156
..... native arsenical,	156
..... arsenicated,	156
..... crude, -	155
..... muriate of,	156
..... red ore of,	156
..... sulphuret of,	156

Antimonial

Page

Antimonial plumoſe ore, 156
..... argentiferous, 156
Antiquities, queries concerning them, - - 177
Antiſeptic powder for preſerving birds - 16, 17
..... ſubſtitute for, 17, 18
Aperient ſaffron of Mars, 71
Apatite, - 106
Apis, - - 6
Aqua fortis -
.... marine - 118
Arcanum duplicatum 99
Argentine, - - 105
..... feltſpar, 125
Argill - - 112
Argillaceous earth, 103, 134
..... iron ſtone 147
..... porphyry 134
..... ſandſtone 134
..... ſchiſtus 116
..... vitriol 69
Argillite, - - 116
Argillo-calcite, 128
..... muriate, 109
Armadilloes, - 189
Armenian bole, 114
Arſenic, 78, 136, 151
..... native, 151
..... oxyde of, 151
..... ſulphuret of, 151
..... red ſulphurized 152
..... miſpickel, 153
..... in water, 79
Arſenical mundick, 152
Aſphaltum, - 166
Aſtronomy of different nations, 179
Atmoſpheric acid, 45, 96
..... air, 83, 92
..... mephitis, 85

Page

Aughtorrha Bhade Shaſtah, 178
Aurora borealis, 185
Auſtrum, - 136

Baikalite, - 112
Baking of birds 20
Banat el-auvi - 189
Barolite, - 107
Baroſelite, - 108
Barotic mephite, 107
..... or ponderous chalk, 107
..... vitriol, 108
Baryte, - 103, 107
Baryto-calcite - 105
Barytum - 137
Baſalt, - 116
Baſaltic hornblende, 115
Baſanite, - - 123
Baſe of marine ſalt, 46, 100
Beans, kinds of, 195
Beetles, - 6, 7
Beryl, - - 119
Bible, manuſcript copies of, 177, 179
Birds, how to deſcribe, xi
.... method of preſerving, 14
..... by
.... the author 13
.... migration of, 189
.... œconomy of, 189, 190
Births and burials, 187
Biſmuth, - 136, 149
..... martial, 150
..... native 149
..... oxyde of, 149
..... ſulphuret of, 149
Bitumen, - 166
..... of Judea, 166
..... maltha, 166
..... petroleum tardifluens, 166

	Page
Bituminous ſubſtances,	165
Black jack -	170
Black lead -	170
Blatta, - -	4
Blende, - -	155
Blood ſtone, -	123
Bole, - -	113
Bolognian ſtone,	108
Blue copperas	75, 144
.... vitriol,	78, 144
Bombylius,	6
Boracic acid, -	96
Boracite, -	112
Borate of ſoda	100
Borax, how procured,	199
Borax, tinkal,	100
Borbonium, -	136
Borrachio, -	196
Botany, queſtions on,	194
Bruchus, .	3
Bupreſtis, -	2, 3
Butter of antimony,	155
Butterflies, where found,	4, 8
..... how to be caught	7
Byrrhus, -	3
Calambac, -	194
Calamine, -	154
Calcareo-ſiliceous ore,	162
..... carbonate,	67, 104
Calcareous earth,	67, 103, 106, 128, 133
..... breccia,	133
..... nitre	64
..... iron ſtone,	147
..... marine ſalt,	65
..... mephite	104
..... muriate,	65
..... nitrate,	64
..... phoſphate,	106
..... ſandſtone,	133
Calcareous ſpar,	67, 104
..... vitriol,	62
Calcedony, - -	121
Calciferous argillite,	130
..... aſbeſtinite,	129
Calcolite, -	159
Calcum, -	137
Caloric, -	11, 84
Calx of arſenic,	78
Cantharis, -	n. 3
Carabus, -	n. 3
Carbonic acid,	45, 85, 96
Carbonates, -	98
Carbonate of alumine,	112
..... ammoniac,	151
..... argill	112
..... baryte,	107
..... copper,	141
..... iron,	71
..... lime,	67, 104
..... magneſia,	60
..... potaſh,	98
..... ſoda,	100
..... zinc,	154
Carbuncle -	119
Carving, art of, -	179
Caſſida, -	n. 3
Catchup, -	184
Catterpillars to feed,	8
Cats-eye, -	122
Cauſtic alkali, -	98
Cicada, -	n. 4
Chalk, - -	67
..... of lead, -	141
..... of zinc, -	154
Chartah Bhade Shaſtah of, Brahmah, - -	178
Chlorite, -	110
Chrome, -	136, 162
Chronology of different nations, -	179

Chryſalis

	Page
Chryſalis of moths and butterflies,	8
Chryſis,	n. 6
Chryſomela,	n. 3
Chryſophraſe	122
Cicindela,	n. 3
Cimex,	n. 4
Circumciſion how uſed,	178
Clay,	112
.... pipe,	112
Clothing, materials of,	128
Coal,	167
Cobalt,	136, 152
.... grey ore of,	153
.... native oxyde of,	153
.... ſulphuret of,	153
.... white ore of,	153
Coccinella,	n. 3
Coin, value of,	180
Coins, impreſſions of,	200
Coleoptera,	n. 2. p. 7
Columbo root,	195
Commerce, queries on,	181
Combuſtion,	164
Common feltſpar,	125
..... hornblende,	115
Compound ſalts,	98
..... ſpar,	105
Conops,	n. 6
Concrete volatile alkali,	101
Copper,	136, 143
..... native,	143
..... oxyde of,	144
..... carbonate of,	144
..... arſeniate of,	144
..... ſulphate of,	145
..... grey ore of,	145
..... white, queries on,	198
..... white ore of,	145
..... muriate of,	144
..... variegated ore of,	145
Copper, ſulphuret of,	145
..... yellow ore of,	145
Copperas,	73
Coral and corallines to collect,	39
Corniſh tin ore,	148
Corundum,	127
Cotton, kinds of,	183
..... how cleaned from ſeeds,	183
Crabs,	6
Cramp or numb-fiſh	191
Cream of lime,	104
Cretaceous acid,	45, 96
Creta,	67
Cretaceous ammoniacal ſalt,	111
..... earth,	67
..... gas,	85
..... magneſia,	60
..... ſoda,	46, 100
..... tartar,	98
Cryſolite,	119
Cryſtals of ſoda,	100
Cubic nitre,	101
.... ſpar,	105
Culinary kitchen ſalt,	53
Curculio,	n. 3
Cuttle fiſh,	193
Cyanite,	114

Davis, captain, his method of preſerving birds,	15
.... obſervation on the ſame,	16
Dermeſtes,	n. 2
Dephlogiſticated air,	82, 92
..... marine acid,	83
Diamond,	116
Diptera claſs,	n. 6. p. 10
Diſeaſes moſt prevalent,	185

	Page
Diſeaſes of manufacturers,	187
Dodecahedral, -	117
Dolomite, -	105
Domeſtication of animals,	186
Drawings of plants, directions for, - -	179
Drawing, ſtate of,	181
Dyeing materials,	181, 182
Ductile metals, -	136
Eagle wood, -	194
Earth, - -	103
..... abſorbent,	104
..... of adamantine ſpar,	103, 127
..... aërated ponderous	107
..... argillaceous	103
..... of bones,	106
..... calcareous	128
..... elementary or ſimple,	103
..... jargon,	103, 127
..... magneſian,	129, 133
..... aërated calcareous,	104
..... ponderous,	103
..... ſiliceous,	103, 132, 134
..... Sidneian,	103, 128
..... ſimple, -	103
..... ſtrontian,	103, 107
..... vitrifiable,	105
..... hornſtone	132
Earthy quartz,	132
.... with actinolite,	132
Echini, or ſea eggs, to preſerve,	21, 193
Efferveſcent barytes,	107
..... calcareous earth,	104
..... mineral fixed alkali, - -	100
..... ſoda,	46
Effluvia of plants,	186
Eggs of birds, parts of zoology, -	23, 189
Egyptian pebble,	123
Elater, - -	n. 3
Electrical rods, -	187
Elephants -	188*
Engliſh ſal volatile,	101
Engravings in ſtone,	180
Ephemera, -	n. 5
Epſom ſalt, -	42, 58
Eudiometer, - -	90
Febrifuge ſalt of Sylvius,	99
Feltſpar, -	124
Ferruginous argillite,	130
..... clays,	131
..... hornſtone,	132
..... limeſtone,	129
..... ſandſtone,	129
..... quartz,	132
Fire damps, -	88
Fiſh to deſcribe -	xii
..... œconomy of,	191
..... to preſerve	20
Fixed air,	42, 45, 85, 92, 96
.... in water,	45
..... alkali, - -	98
..... ſal ammoniac,	65
Flint, - - -	122
Flowers of arſenic,	78
Fluate of lime, -	105
Fluor ſpar, -	105
..... volatile alkali,	101
Fluoric acid gas,	82
Food of different nations	185
Forficula, -	n. 4
Foſſils, directions in collecting,	93
Fragile metals -	33
Frogs to preſerve,	21
Fuci, - -	192
Fullers earth, -	113

	Page
Fulminating gold,	137
Garil,	104
Garnet,	119
Gas, deſcription of,	81
.... dephlogiſticated muriatic acid	157
.... hepatic	165
.... ſylveſtre	45, 96
.... ſuroxygenated muriatic,	157
Gems, whence produced,	199
Glauber's ſalt,	48, 101
Gold,	136, 137
.... grey ore of,	138
.... native	138
.... white ore of,	138, 163
.... graphic of Offenbanya	163
Goſlard vitriol,	77
Green copperas,	147
.... vitriol,	147
Gryllus,	n. 4
Gum ammoniac,	197
.... myrrh,	197
.... rubrum aſtringens from Gambia,	197
Gypſum,	43, 62, 106, 129
Gyrinus,	n. 3
Hæmatites,	31
Heavy ſpar,	108
Heliotropium,	123
Hemerobius,	n. 5
Hemiptera claſs,	n. 4. p. 7
Hepatic gas,	83, 165
Hexagonal,	107
Hippoboſca,	n. 6
Hiſter,	n. 2
Holothuriæ,	193
Homogeneous earth,	103
Honeyſtone,	168
Hornſlate,	123
Hornſtone,	123
Hornblende baſaltic,	115
..... Labradore	115
..... with garnet,	130
.... talc or mica,	130
..... ſlate with quartz,	131
..... ſchiſtoſe.	115
Humming birds,	190
Hyacinth,	119
Hyalite,	122
Hydrogen gas,	154, 164
..... or inflammable air,	87, 92
..... pure,	164
..... ſulphurated,	165
Hydrophanes,	122
Hymenoptera claſs,	n. 5. p. 10
Ichneumon,	n. 5
Icoſahedrons,	118
Jacinth,	119
Jade,	111
Jackal,	189
Jargon,	127
..... earth,	101, 127
Jaſper,	123
Idols, hiſtories of,	178, 179
Jet,	167
Ignis fatuus	88
Impure air,	85
Indigo, manufactory of,	183
Indian nut,	195
Indurated talc,	109
Inflammable gas,	82, 83, 92
Inflammation,	164
Inſcriptions	164
Inſects, method of preſerving,	1
..... method of killing,	8, 9

	Page
Infects, method of securing from vermin,	11, 12
..... to describe,	xiii
Journal of occurrences,	xvi
Iron and it's ores,	31, 136, 145
.... Argillaceous ore of,	146
.... grey ore of,	146
.... hæmatite,	146
.... native,	146
.... sulphuret of,	147
.... spathose ore of,	146
.... stone,	147
.... sulphate of,	147
Kali vitriolatum,	50
Kaoutchuck, an elastic gum,	196
Killas,	116
Krag,	116
Kramer's antiseptic powder for preserving birds,	11, 16
Kuckahn's method of preserving birds,	15, 17
..... observations on,	15, 17
Kupfer nickel,	150
Labrador feltspar,	125
..... hornblende,	115
Lac lunæ,	112
Lama's sacred books,	178
Lampyris,	n. 3
Lapis calaminaris,	154
.... lazuli,	126
.... lydius,	126
.... ollaris,	109
Latent heat,	n. 84
Lava,	173
.... vitreous,	173
Lead,	26, 136, 140
.... amalgam,	141
.... antimonial ore of,	142
Lead black,	170
.... black spar of,	142
.... brown spar of,	142
.... carbonate of,	141
.... chalk of,	141
.... galæna, sulphuret of lead, or sulphurized ore of,	142
.... mephite of,	141
.... native,	141
.... native glass of,	142
.... phosphorated ore of,	143
.... pyritical ore of,	142
.... red ore of,	141
.... spathose,	141
.... transparent,	141
.... white opaque ore of,	141
.... yellow ore of,	143
Learning, queries respecting,	177
Leman's method of preserving birds,	n. 17
Leopards,	189
Lepidolite,	114
Lepidoptera class,	n. 4. p. 8
Leptura,	n. 3
Libellula,	n. 5
Lignum Rhodium,	194
Lime,	103
.... of arsenic,	78
Limestone,	104
..... with argillite,	129
Linnæus's method of preserving birds,	16
..... seeds,	26
Lithomarga,	113
Liver of stone,	108
..... sulphur in water,	43
Lizards to preserve,	18
Lobsters,	6
Lucanus,	n. 4
Madrepores,	4

	Page
Magnesia,	60, 109
Magnesian,	14, 15
..... carbonate,	60
..... chalk,	60
..... earth,	60, 103, 133
..... muriate,	43, 59
..... nitre,	42. 58
..... vitriol,	55
Magnesium,	137
Maltha,	166
Manganese,	136, 157
..... black calciform,	158
..... native,	157
..... oxyde of,	157
..... siliceous ore of,	158
..... white oxydated,	157
Manganesean wolfram,	159
Mantis,	n. 4
Manufactures, queries upon,	181
Marble,	104
Marine alkali,	46
Marl,	128
Marriage ceremonies,	179
Martial chalk,	71
..... mephite,	71
..... muriatic spar,	109
..... vitriol,	73, 147
Medals, impressions of,	201
Medicinal waters, method of analyzing	41
Meloa,	n. 3
Menachanite,	160
..... native,	160
Mendæans, sacred books of,	178
Mephite of lead,	141
..... magnesia,	60
..... potash,	98
..... zinc,	154
Mephitic acid,	45, 85, 96
..... tartar,	98
Mercury or quicksilver,	116
	138
Metallic substances in water,	73
Metals to collect,	93, 200
..... the knowledge of their uses,	199
Meteorological observations recommended,	185
Mica or glimmers,	115
Millepores,	4
Mineral alkali,	46, 100
..... elastic gum,	167
..... pitch,	166
..... tallow,	169
..... water to analyze,	41
Mineralogy, queries upon,	198
Mines, method of working,	198
Mispickel,	152
Mollusca,	192
Molybdate of lead,	141
Moonstone,	124
Mother water of nitre,	64
..... marine salt,	65
Moths, where found,	4
..... how caught,	8
..... method of killing,	8
Mountain potash,	166
..... soap,	113
Mullen,	131
..... with asbestinate,	131
..... siliceous schistus,	133
..... trap,	131
Muriate of ammoniac,	101
..... lime,	65
..... magnesia,	43, 59
..... oxygenated of potash,	99
..... potash,	99
..... soda,	53
Muriated ammoniac,	101
Muriates,	99, 144

Muriatic

	Page
Muriatic acid gas,	82
..... earth,	60
Muri-calcite,	105
Muſca,	n. 6
Muſcovy glaſs,	115
Mutilla,	n. 6
Myrmillion,	n. 5
Myrobalans,	184
Myrrh, how produced,	197
Mytilus margaritifer,	193
Nagyag, yellow ore of,	163
..... foliated grey gold of,	163
Naptha,	165
Natrum,	46
Natrum vitriolatum,	48, 100
Nautilus,	192
Necydalis,	n. 3
Nepa,	n. 4
Neſts of birds,	22
Net for catching water inſects,	3
..... inſects when flying,	9
Neuroptera claſs,	n. 5 p. 10
Nickel,	136, 150
..... native,	150
..... calx or oxyde of,	150
..... kupfer,	150
Nitrate of ammoniac,	102
.... lime,	64
.... magneſia,	58
.... potaſh,	99
.... ſoda,	101
Nitrates,	99
Nitrate of potaſh,	99
Nitre or ſaltpetre,	99
.... fixed by itſelf,	98
.... of magneſia,	58
.... .. potaſh,	52
Notonecta,	4
Novaculite,	116
Nyctanthus,	194
Obſidian,	121
Octohedrons,	116
Oeſtrus,	n. 6
Oil of vitriol,	97
Olivia,	119
Opal,	122
Orpiment, yellow,	151
Ounces,	189
Oxyde of arſenic,	78
Oxygen, definition of,	82
..... gas,	84
Oxygenated arſenic,	78
..... muriatic acid gas,	83
Palm trees, kinds of,	195
Panthers,	189
Papilio,	n. 4
Parthenum,	136
Peacocks, wild,	191
Pearl ſpar,	104
Pearls, formation of,	193
Peas,	195
Pedra de cobra,	186
Pehlvi,	178
Petroleum,	165
Petroſilex,	123
Prehnite,	126
Phalæna,	n. 4
Phaſeoli, kinds of,	195
Phlogiſticated air,	85, 92
..... gas,	85
..... vitriolic acid,	97
Phlogiſton of Kirwan,	87
Phoſphate of alumine,	112
..... baryte,	108
..... lime,	106
Phoſphoric gas,	83
..... ſpar,	105
Phoſphorated hydrogenated gas,	83
Phrygamia,	n. 5
Pipe clay,	112

	Page
Pitchſtone,	122
..... with opal,	133
Plants to preſerve and tranſport,	23
.... to deſcribe and draw,	xi. xv
.... to dry,	35
.... to take impreſſions of,	39
Plaſter of Paris,	62, 106
.... Paris in water,	62
..... ſtone,	62
Platina,	137
Plumbago,	170
..... common,	170
Poiſons, effects of,	186
Polite arts, queries upon,	177
Polygamy, how employed,	186
Polypes,	193
Ponderous earth,	103, 107
..... clay,	113
..... ſpar,	107, 108
Porcelain,	113
Potaſh,	98
Potſtone,	109
..... Porphyry,	133
Praſe,	121
Pterocarpus,	196
Ptinus,	n. 3
Pudding ſtone	135
Pulſe, kinds of,	195
Pumice,	172
Purple of the ancients,	193
Puzzolana,	171

	Page
Quadrupeds to deſcribe,	xi
.... catch,	182
..... œconomy of,	188
Quartz,	121, 132
Quickſilver,	136, 138
..... analyzed,	139
..... cupreous,	139
..... hepatic ore of,	139
..... native,	139

	Page
Raphidia,	n. 5
Rattleſnakes, enquiries concerning,	192
Ravens, Indian,	190
Realgar,	151
Religious rites, queries on,	177
Reptiles to preſerve,	21
Reſplendent hornblende	115
Rhomboidal nitre,	101
Rice, kinds of,	196
Riuté, ſacred book,	178
Rochel,	194
Rock oil,	165
Roſe ſpar,	127
Rubellite,	120
Ruby,	118
Ruby arſenic,	152
Ruddle ſtone	134
Ruſt of iron,	71

Sal ammoniac,	101
.. catharticus,	48
.. de duobus	80
.. gemmæ,	53
.. marinum	53
.. martis,	73
.. mirabile,	48
.. polychreſtum,	50
.. polychreſt of Glaſer,	99
.. vitrioli,	77
Salt, common,	53
.... petre,	42, 99
.... how manufactured in India,	199
.... of Sedlitz,	55
.... acid and neuter,	45
.... ſimple,	96
.... in water to detect,	42
Salmiac,	101
Sand,	171
Sappare,	114
Sapphire,	117

Scarabæus,

	Page
Scarabæus, chafer,	n. 2
Scheele, . 136,	158
Scheifer ſpar, .	105
Schiller ſpar, .	115
Schiſtoſe hornblende,	115
..... mica, .	134
Schiſtoſe talc, .	109
Schorl, . .	120
Schorlite, . .	120
Scoriæ, .	174
Sea cow, .	191
Sea eggs, . 19,	193
... fowl, how to preſerve,	20
... weeds to be examined,	192
Seaſons, ſucceſſion of,	185
Sedative ſalt, .	96
Seeds to preſerve and tranſport, xv. 23,	26
.... when to be ſown,	27
Selenite, . .	106
..... in water,	62
Semi-opal, .	122
Sepia, . .	193
Sepulchral rites, .	179
Serpentine, . .	110
..... with hornblende,	129
.... porphyry,	133
Shells to deſcribe,	xiv
Ship-building, .	183
Sidero-calcite, .	104
Sidneia, . .	128
Sidneian earth, 103,	128
Sienite, .	134
Silex, . . 103,	116
Siliceous breccia,	135
..... earth, 103,	134
..... iron ſtone,	147
..... limeſtone,	128
..... porphyry,	135
..... ſpar, .	126
..... ſandſtone,	135
..... ſchiſtus,	123
.... argillite,	132

	Page
Siliceous ſchiſtus, with limeſtone,	132
.... mullen,	133
.... opal,	133
Silici-murite, .	109
Siliciferous argillite,	130
..... potſtone,	129
..... trap,	131
Silk grey in China,	192
Silpha, . .	n. 2
Silver, . 136,	138
..... arſenical,	140
..... brittle, vitreous,	140
..... horn, .	140
..... native, .	140
..... red ore of,	140
..... vitreons, .	140
..... white ore of,	140
Slaggs, . .	174
Smalts, . .	153
Smaragdite, .	118
Soda, . .	46
.... vitriolata, .	48
Solid air of Hali,	85
Soya, . .	184
Snakes to deſcribe,	xii. 21
..... poiſon of,	186
Sparry fluor, .	105
Spathoſe lead, .	141
..... tin ore, .	148
Sphex, . .	n. 6
Sphinx, .	n. 4
Spiritus Sylveſtris,	45
Sponges to collect, 39,	193
Steatites, . .	110
..... with argill,	129
Strawberry ſeeds to preſerve,	25
Stream tin, .	148
Strontian earth, 103,	107
Strontionite, .	107
Sublimed white oxyde of arſenic, . .	78
Sulphate of alumine,	69

	Page
Sulphate of ammoniac,	102
..... baryte, .	108
..... copper, 75,	145
..... iron, . .	73
..... of lead, .	142
..... lime, 62,	106
..... magnesia,	55
..... potash, 50,	99
..... soda, 48,	100
..... zinc, 75,	155
Sulphites, . .	98
Sulphur, . .	169
..... common,	170
..... volcanic,	170
Sulphures, . .	97
Sulphuret of antimony,	156
..... arsenic,	151
..... bismuth,	149
..... copper,	145
..... tin, .	148
Sulphuric acid, .	97
Sulphurised bismuth,	149
..... tin,	148
..... zinc,	155
Sylvanite, . .	161
Sympathetic ink,	153
Swallows, when first seen,	190
Tabanus, .	n. 6
Tafa, . .	172
Talc, . .	109
Talcite, . .	109
Talcose argillite,	130
Tallow mineral,	169
Tatack, a species of grass,	195
Tea seeds to preserve,	27
Tek, what species of wood,	195
Tellurium, . 106,	162
Tenebrio, . .	n. 3
Terra lemnia, .	114
Terrass, . .	172
Thumerstein, .	120
Tigers, . .	189

	Page
Tin, . . 136,	148
Tin, brown tin-stone,	148
..... pyrites, .	148
..... stream, or Cornish ore of,	148
..... sulphuret of,	148
Tipula, . .	n. 6
Titanite, . 136,	161
Topaz, . .	117
Tortoises, to describe,	xii
..... preserve,	21
Tourmalin, .	120
Trade, queries respecting,	181
Trap, . .	116
..... with hornblende,	131
..... krag,	131
..... mullen,	131
Trass, . .	172
Tripoli, . .	114
Tsai, an Indian plant for dyeing, . .	196
Tunstate, .	158
..... of lime,	158
Tungstein, . 136,	158
Uranite, . 136,	159
..... calciform of,	159
..... carbonate of,	159
..... oxyde of,	159
..... spathose,	159
..... sulphuret of,	160
Uranium, . .	159
Vedam, a sacred book of the Bramins, .	178
Vegetable alkali,	98
..... fixed alkali,	98
Venetian talc, .	109
Vespa, . .	n. 6
Vesuvian, .	120
Violet schorl, .	120
Vital air, . .	82
Vitiated air, .	85

	Page
Vitreous ſpar,	105
Vitriolic acid,	97
Vitriol of copper,	75
..... Cyprus,	144
..... goſlar,	155
..... green,	147
..... of iron,	73, 147
..... lime,	106
..... potaſh,	99
..... ſoda,	100
..... Venus,	75, 144
..... zinc,	97, 155
Vitriolum viride,	73
Volatile alkaline gas,	83
..... narcotic ſalt of vitriol,	96
..... ſpirit of ſal ammoniac,	101
Wakken,	116
Waſps,	5
White arſenic,	78
..... copperas,	77, 155
..... oxyde of arſenic,	78
White vitriol,	77, 155
Wolfram,	41, 136, 156
Wood tin,	148
Yellow orpiment,	151
Zaffer for painting,	183
Zend,	178
Zeolite,	126
Zinc,	136, 154
..... aërated,	154
..... blende or ſulphurized,	155
..... calx of,	154
..... carbonate of,	154
..... chalk of,	154
..... mephite of,	154
..... native oxyde of,	154
..... vitriol of,	155
..... ſulphate of,	155
Zirgon,	127
..... earth,	127
Zoology, queries concerning,	188
Zuſa,	104

THE END.